

Anti-Kinesin 1 / UKHC Antibody (C-term), Biotinylated

Catalog # AF4276a

Product Information

Application	WB, IHC, E
Primary Accession	P33176
Other Accession	3799 , NP_004512.1 , 16573 , 117550
Reactivity	Human, Dog
Predicted	Human, Mouse, Rat, Dog
Calculated MW	109685

Additional Information

Gene ID	3799
Other Names	cytoskeletal; ERM; acetylprotein
Dilution	WB~~1:1000 IHC~~1:100~500 E~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Kinesin 1 / UKHC Antibody (C-term), Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	KIF5B (HGNC:6324)
Synonyms	KNS, KNS1
Function	Microtubule-dependent motor required for normal distribution of mitochondria and lysosomes. Can induce formation of neurite-like membrane protrusions in non-neuronal cells in a ZFYVE27-dependent manner (By similarity). Regulates centrosome and nuclear positioning during mitotic entry. During the G2 phase of the cell cycle in a BICD2- dependent manner, antagonizes dynein function and drives the separation of nuclei and centrosomes (PubMed: 20386726). Required for anterograde axonal transportation of MAPK8IP3/JIP3 which is essential for MAPK8IP3/JIP3 function in axon elongation (By similarity). Through binding with PLEKHM2 and ARL8B, directs lysosome movement toward microtubule plus ends (Probable). Involved in NK cell-mediated cytotoxicity. Drives the polarization of cytolytic granules and microtubule-organizing centers (MTOCs) toward the immune synapse between effector NK lymphocytes and target cells (PubMed: 24088571).

Cellular Location

Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q2PQA9}. Cytolytic granule membrane. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side Note=Uniformly distributed between soma and neurites in hippocampal neurons. {ECO:0000250|UniProtKB:Q2PQA9}

Images



Biotinylated antibody (1 µg/ml) staining of HeLa lysate (35 µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.