

# Goat Anti-UBID4 Antibody (aa89-103) (internal region)

Catalog # AF4297a

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q92785</a>
Other Accession	<a href="#">NP_006259.1</a> , <a href="#">361711</a> , <a href="#">19708</a> , <a href="#">5977</a>
Reactivity	Human
Predicted	Human, Mouse, Rat, Pig, Dog
Host	Goat
Isotype	IgG
Calculated MW	44155

## Additional Information

---

Gene ID	5977
Other Names	Zinc finger protein ubi-d4, Apoptosis response zinc finger protein, BRG1-associated factor 45D, BAF45D, D4, zinc and double PHD fingers family 2, Protein requiem, DPF2, BAF45D, REQ, UBID4
Dilution	WB~~1:1000 E~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat Anti-UBID4 Antibody (aa89-103) (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.


## Protein Information

---

Name	DPF2
Synonyms	BAF45D, REQ, UBID4
Function	Plays an active role in transcriptional regulation by binding modified histones H3 and H4 (PubMed: <a href="#">27775714</a> , PubMed: <a href="#">28533407</a> ). Is a negative regulator of myeloid differentiation of hematopoietic progenitor cells (PubMed: <a href="#">28533407</a> ). Might also have a role in the development and maturation of lymphoid cells (By similarity). Involved in the regulation of non-canonical NF-kappa-B pathway (PubMed: <a href="#">20460684</a> ).
Cellular Location	Nucleus. Cytoplasm
Tissue Location	Ubiquitous.

## Images

---



AF4297a (1  $\mu$ g/ml) staining of Jurkat (A) and K562 (B) lysate (35  $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.