

Goat Anti-UBID4 Antibody (aa89-103) (internal region)

Catalog # AF4297a

Product Information

Application	WB, E
Primary Accession	Q92785
Other Accession	NP_006259.1 , 361711 , 19708 , 5977
Reactivity	Human
Predicted	Human, Mouse, Rat, Pig, Dog
Host	Goat
Isotype	IgG
Calculated MW	44155


Additional Information

Gene ID	5977
Other Names	Zinc finger protein ubi-d4, Apoptosis response zinc finger protein, BRG1-associated factor 45D, BAF45D, D4, zinc and double PHD fingers family 2, Protein requiem, DPF2, BAF45D, REQ, UBID4
Dilution	WB~~1:1000 E~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat Anti-UBID4 Antibody (aa89-103) (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DPF2
Synonyms	BAF45D, REQ, UBID4
Function	Plays an active role in transcriptional regulation by binding modified histones H3 and H4 (PubMed: 27775714 , PubMed: 28533407). Is a negative regulator of myeloid differentiation of hematopoietic progenitor cells (PubMed: 28533407). Might also have a role in the development and maturation of lymphoid cells (By similarity). Involved in the regulation of non-canonical NF-kappa-B pathway (PubMed: 20460684).
Cellular Location	Nucleus. Cytoplasm
Tissue Location	Ubiquitous.

Images



AF4297a (1 μ g/ml) staining of Jurkat (A) and K562 (B) lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.