

# Goat Anti-CDK9 Antibody

Peptide-affinity purified goat antibody

Catalog # AF4321a


## Product Information

|                        |                                     |
|------------------------|-------------------------------------|
| <b>Application</b>     | WB, E                               |
| <b>Other Accession</b> | <a href="#">NP_001252.1</a>         |
| <b>Reactivity</b>      | Human, Mouse, Rat, Pig, Dog, Bovine |
| <b>Host</b>            | Goat                                |

## Additional Information

|                    |                                                                                                                                                                                                                                                                  |
|--------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Other Names</b> | CDK9, cyclin-dependent kinase 9, C-2k, CDC2L4,, CTK1, PITALRE, TAK, CDC2-related kinase,, cell division cycle 2-like protein kinase 4, cell division protein kinase 9,, serine/threonine protein kinase PITALRE, tat-associated kinase complex catalytic subunit |
| <b>Dilution</b>    | WB~~1:1000 E~~N/A                                                                                                                                                                                                                                                |
| <b>Storage</b>     | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.                                                                                                                         |
| <b>Precautions</b> | Goat Anti-CDK9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.                                                                                                                                                        |

## Images



AF4321a (0.5 µg/ml) staining of NIH3T3 nuclear lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.