

## Goat anti-IL6R / CD126 (isoform 1) Antibody

Peptide-affinity purified goat antibody Catalog # AF4338a

## **Product Information**

**Application** Pep-ELISA **Primary Accession** P08887 **Other Accession** NP 000556.1 Reactivity Human Host Goat Clonality Polyclonal **Clone Names** IL6R **Calculated MW** 51548

## **Additional Information**

**Gene ID** 3570

Other Names IL6R; interleukin 6 receptor; HGNC:6019; CD126; IL-6R-1; IL-6R-alpha; IL6RA;

MGC104991; CD126 antigen; interleukin 6 receptor alpha subunit

**Dilution** Pep-ELISA~~N/A

**Format** Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5%

bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and

thawing.

**Immunogen** This antibody is expected to recognise isoform 1 only (NP\_000556.1)

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** Goat anti-IL6R / CD126 (isoform 1) Antibody is for research use only and not

for use in diagnostic or therapeutic procedures.

## **Protein Information**

Name IL6R ( HGNC:6019)

**Function** Part of the receptor for interleukin 6. Binds to IL6 with low affinity, but does

not transduce a signal (PubMed:<u>28265003</u>). Signal activation necessitate an association with IL6ST. Activation leads to the regulation of the immune response, acute-phase reactions and hematopoiesis (PubMed:<u>30995492</u>, PubMed:<u>31235509</u>). The interaction with membrane-bound IL6R and IL6ST stimulates 'classic signaling', the restricted expression of the IL6R limits classic IL6 signaling to only a few tissues such as the liver and some cells of

the immune system. Whereas the binding of IL6 and soluble IL6R to IL6ST stimulates 'trans- signaling'. Alternatively, 'cluster signaling' occurs when membrane- bound IL6:IL6R complexes on transmitter cells activate IL6ST receptors on neighboring receiver cells (Probable).

Cellular Location [Isoform 1]: Cell membrane {ECO:0000250|UniProtKB:P22272}; Single-pass

type I membrane protein [Soluble interleukin-6 receptor subunit alpha]:

Secreted

**Tissue Location** [Isoform 2]: Expressed in peripheral blood mononuclear cells and weakly

found in urine and serum. 1%-20% of the total sIL6R in plasma is generated

by alternative splicing (PubMed:28060820).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.