

Goat anti-TXNDC1 / TMX, biotinylated Antibody

Peptide-affinity purified goat antibody

Catalog # AF4350a

Product Information

Application	WB, Pep-ELISA
Primary Accession	Q9H3N1
Other Accession	NP_110382.3
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Clone Names	TMX1
Calculated MW	31791

Additional Information

Gene ID	81542
Other Names	TMX1; thioredoxin-related transmembrane protein 1; PDIA11; TMX; TXNDC; TXNDC1; protein disulfide isomerase family A, member 11; thioredoxin domain containing 1; thioredoxin domain-containing protein 1; transmembrane Trx-related protein
Dilution	WB~~1:1000 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-TXNDC1 / TMX, biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TMX1 {ECO:0000303 PubMed:37648867, ECO:0000312 HGNC:HGNC:15487}
Function	Thioredoxin domain-containing protein that participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions (PubMed: 11152479 , PubMed: 37648867). Acts as a key inhibitor of the alternative triglyceride biosynthesis pathway by inhibiting the activity of TMEM68/DIESL at the endoplasmic reticulum, thereby restricting accumulation of triacylglycerol (PubMed: 37648867). The alternative triglyceride biosynthesis pathway

mediates formation of triacylglycerol from diacylglycerol and membrane phospholipids (PubMed:[37648867](#)). Acts as a protein disulfide isomerase by catalyzing formation or reduction of disulfide bonds (PubMed:[22228764](#), PubMed:[29932915](#)). Specifically mediates formation of disulfide bonds of transmembrane proteins at the endoplasmic reticulum membrane (PubMed:[22228764](#)). Involved in endoplasmic reticulum-associated degradation (ERAD) via its protein disulfide isomerase activity by acting on folding-defective polypeptides at the endoplasmic reticulum membrane (PubMed:[29932915](#)). Acts as a negative regulator of platelet aggregation following secretion in the extracellular space (PubMed:[30425049](#)). Acts as a regulator of endoplasmic reticulum- mitochondria contact sites via its ability to regulate redox signals (PubMed:[27502484](#), PubMed:[31304984](#)). Regulates endoplasmic reticulum- mitochondria Ca(2+) flux (PubMed:[27502484](#)).

Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein. Mitochondrion membrane; Single-pass type I membrane protein. Secreted. Note=Predominantly found in the endoplasmic reticulum (PubMed:11152479). Secreted in the extracellular space following thrombin stimulation (PubMed:30425049). Localizes to mitochondria-associated endoplasmic reticulum membrane (MAM); palmitoylation is required for MAM localization (PubMed:22045338, PubMed:27502484, PubMed:31304984).

Tissue Location

Ubiquitous (PubMed:11152479). Highly expressed in kidney, liver, placenta and lung (PubMed:11152479)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.