

Goat anti-CBX5 / HP1-Alpha, biotinylated Antibody

Peptide-affinity purified goat antibody

Catalog # AF4353a

Product Information

Application	WB, Pep-ELISA
Primary Accession	P45973
Other Accession	NP_036249.1
Reactivity	Human, Mouse, Dog
Host	Goat
Clonality	Polyclonal
Clone Names	CBX5
Calculated MW	22225

Additional Information

Gene ID	23468
Other Names	CBX5; chromobox homolog 5; HEL25; HP1; HP1A; HP1 alpha homolog; HP1-ALPHA; HP1Hs alpha; antigen p25; chromobox homolog 5 (HP1 alpha homolog, Drosophila); epididymis luminal protein 25; heterochromatin protein 1 homolog alpha; heterochromatin protein 1-alp
Dilution	WB~~1:1000 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Immunogen	Reported variants represent identical protein: NP_036249.1, NP_001120793.1, NP_001120794.1
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-CBX5 / HP1-Alpha, biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CBX5
Synonyms	HP1A
Function	Component of heterochromatin that recognizes and binds histone H3 tails methylated at 'Lys-9' (H3K9me), leading to epigenetic repression. In contrast,

it is excluded from chromatin when 'Tyr-41' of histone H3 is phosphorylated (H3Y41ph) (PubMed:[19783980](#)). May contribute to the association of heterochromatin with the inner nuclear membrane by interactions with the lamin-B receptor (LBR) (PubMed:[19783980](#)). Involved in the formation of kinetochore through interaction with the MIS12 complex subunit NSL1 (PubMed:[19783980](#), PubMed:[20231385](#)). Required for the formation of the inner centromere (PubMed:[20231385](#)).

Cellular Location

Nucleus. Chromosome. Chromosome, centromere. Note=Colocalizes with HNRNPU in the nucleus (PubMed:19617346). Component of centromeric and pericentromeric heterochromatin. Associates with chromosomes during mitosis. Associates specifically with chromatin during metaphase and anaphase (PubMed:19617346). Localizes to sites of DNA damage (PubMed:28977666)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.