

Goat anti-UBE2I, Biotinylated Antibody

Peptide-affinity purified goat antibody Catalog # AF4362a

Product Information

Application	WB, IHC, ICC, Pep-ELISA
Primary Accession	<u>P63279</u>
Other Accession	<u>NP_003336.1</u>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Clone Names	UBE2I
Calculated MW	18007

Additional Information

Gene ID	7329
Other Names	UBE2I; ubiquitin-conjugating enzyme E2I; C358B7.1; P18; UBC9; SUMO-1-protein ligase; SUMO-protein ligase; ubiquitin carrier protein 9; ubiquitin carrier protein I; ubiquitin conjugating enzyme 9; ubiquitin-conjugating enzyme E2I (UBC9 homolog, yeast); ubi
Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Immunogen	Reported variants represent identical protein: NP_919237.1, NP_003336.1, NP_919236.1, NP_919235.1
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-UBE2I, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	UBE2I
Synonyms	UBC9, UBCE9
Function	Accepts the ubiquitin-like proteins SUMO1, SUMO2, SUMO3, SUMO4 and SUMO1P1/SUMO5 from the UBLE1A-UBLE1B E1 complex and catalyzes their

	covalent attachment to other proteins with the help of an E3 ligase such as RANBP2, CBX4 and ZNF451. Can catalyze the formation of poly-SUMO chains. Necessary for sumoylation of FOXL2 and KAT5. Essential for nuclear architecture and chromosome segregation. Sumoylates p53/TP53 at 'Lys-386'. Mediates sumoylation of ERCC6 which is essential for its transcription-coupled nucleotide excision repair activity (PubMed: <u>26620705</u>).
Cellular Location	Nucleus. Cytoplasm Cytoplasm, perinuclear region Note=Mainly nuclear (By similarity). In spermatocytes, localizes in synaptonemal complexes (PubMed:8610150). Recruited by BCL11A into the nuclear body (By similarity). {ECO:0000250 UniProtKB:P63280, ECO:0000269 PubMed:8610150}
Tissue Location	Expressed in heart, skeletal muscle, pancreas, kidney, liver, lung, placenta and brain. Also expressed in testis and thymus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.