

Goat anti-ACADM, Biotinylated Antibody

Peptide-affinity purified goat antibody

Catalog # AF4364a

Product Information

Application	WB, IHC, Pep-ELISA
Primary Accession	P11310
Other Accession	NP_000007.1 , NP_001120800.1 , NP_001272971.1 , NP_001272972.1 , NP_001272973.1
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Clone Names	ACADM
Calculated MW	46588

Additional Information

Gene ID	34
Other Names	ACADM; acyl-CoA dehydrogenase, C-4 to C-12 straight chain; ACAD1; MCAD; MCADH; acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight chain
Dilution	WB~~1:1000 IHC~~1:100~500 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Immunogen	This antibody is expected to recognise both reported isoforms (NP_000007.1; NP_001120800.1).
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-ACADM, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ACADM (HGNC:89)
Function	Medium-chain specific acyl-CoA dehydrogenase is one of the acyl-CoA dehydrogenases that catalyze the first step of mitochondrial fatty acid beta-oxidation, an aerobic process breaking down fatty acids into acetyl-CoA and allowing the production of energy from fats (PubMed: 1970566 , PubMed: 21237683 , PubMed: 2251268 , PubMed: 8823175). The first step of

fatty acid beta-oxidation consists in the removal of one hydrogen from C-2 and C-3 of the straight-chain fatty acyl-CoA thioester, resulting in the formation of trans-2-enoyl-CoA (PubMed:[2251268](#)). Electron transfer flavoprotein (ETF) is the electron acceptor that transfers electrons to the main mitochondrial respiratory chain via ETF-ubiquinone oxidoreductase (ETF dehydrogenase) (PubMed:[15159392](#), PubMed:[25416781](#)). Among the different mitochondrial acyl-CoA dehydrogenases, medium-chain specific acyl-CoA dehydrogenase acts specifically on acyl-CoAs with saturated 6 to 12 carbons long primary chains (PubMed:[1970566](#), PubMed:[21237683](#), PubMed:[2251268](#), PubMed:[8823175](#)).

Cellular Location

Mitochondrion matrix

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.