

Goat anti-ITGA3 (isoform a) Antibody

Peptide-affinity purified goat antibody

Catalog # AF4390a

Product Information

Application	WB, Pep-ELISA
Primary Accession	P26006
Other Accession	NP_002195.1
Reactivity	Human, Mouse, Pig, Bovine
Host	Goat
Clonality	Polyclonal
Clone Names	ITGA3
Calculated MW	116612

Additional Information

Gene ID	3675
Other Names	ITGA3; integrin, alpha 3 (antigen CD49C, alpha 3 subunit of VLA-3 receptor); CD49C; GAP-B3; GAPB3; ILNEB; MSK18; VCA-2; VL3A; VLA3a; CD49 antigen-like family member C; FRP-2; VLA-3 subunit alpha; antigen identified by monoclonal antibody J143; galactoprot
Dilution	WB~~1:1000 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Immunogen	This antibody is expected to recognize reported isoform a (NP_002195.1).
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-ITGA3 (isoform a) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ITGA3
Synonyms	MSK18
Function	Integrin alpha-3/beta-1 is a receptor for fibronectin, laminin, collagen, epiligrin, thrombospondin and CSPG4. Integrin alpha- 3/beta-1 provides a docking site for FAP (seprase) at invadopodia plasma membranes in a

collagen-dependent manner and hence may participate in the adhesion, formation of invadopodia and matrix degradation processes, promoting cell invasion. Alpha-3/beta-1 may mediate with LGALS3 the stimulation by CSPG4 of endothelial cells migration.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell membrane; Lipid-anchor. Cell projection, invadopodium membrane; Single-pass type I membrane protein. Cell projection, filopodium membrane; Single-pass type I membrane protein. Note=Enriched preferentially at invadopodia, cell membrane protrusions that correspond to sites of cell invasion, in a collagen-dependent manner.

Tissue Location

Isoform 1 is widely expressed. Isoform 2 is expressed in brain and heart. In brain, both isoforms are exclusively expressed on vascular smooth muscle cells, whereas in heart isoform 1 is strongly expressed on vascular smooth muscle cells, isoform 2 is detected only on endothelial vein cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.