

# Goat anti-LAG3 / CD223 Antibody

Peptide-affinity purified goat antibody

Catalog # AF4463a

## Product Information

---

Application	WB, Pep-ELISA
Primary Accession	<a href="#">P18627</a>
Other Accession	<a href="#">NP_002277.4</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Clone Names	LAG3
Calculated MW	57449

## Additional Information

---

Gene ID	3902
Other Names	LAG3; lymphocyte activating 3; CD223; lymphocyte-activation gene 3
Dilution	WB~~1:1000 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Immunogen	This antibody is expected to recognize the extra-cellular domain.
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-LAG3 / CD223 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	LAG3 ( <a href="#">HGNC:6476</a> )
Synonyms	FDC
Function	Lymphocyte activation gene 3 protein: Inhibitory receptor on antigen activated T-cells (PubMed: <a href="#">20421648</a> , PubMed: <a href="#">7805750</a> , PubMed: <a href="#">8647185</a> ). Delivers inhibitory signals upon binding to ligands, such as FGL1 (By similarity). FGL1 constitutes a major ligand of LAG3 and is responsible for LAG3 T-cell inhibitory function (By similarity). Following TCR engagement, LAG3 associates with CD3-TCR in the immunological synapse and directly

inhibits T-cell activation (By similarity). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1, possibly by acting as a coreceptor for PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed:[20421648](#), PubMed:[7805750](#), PubMed:[8647185](#)). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity). Binds MHC class II (MHC-II); the precise role of MHC-II-binding is however unclear (PubMed:[8647185](#)).

**Cellular Location**

[Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein

**Tissue Location**

Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.