10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Goat anti-ARX Antibody

Peptide-affinity purified goat antibody
Catalog # AF4468a

Product Information

Application	WB, Pep-ELISA
Primary Accession	Q96QS3
Other Accession	NP_620689.1
Reactivity	Human, Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Clone Names	ARX
Calculated MW	58160

Additional Information

Gene ID	170302
Other Names	ARX; aristaless related homeobox; ISSX; MRX29; MRX32; MRX33; MRX36; MRX38; MRX43; MRX54; MRXS1; PRTS; Infantile spasm syndrome, X-linked; aristaless-related homeobox, X-linked
Dilution	WB~~1:1000 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-ARX Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ARX {ECO:0000303|PubMed:12359145, ECO:0000312|HGNC:HGNC:18060}
Function	DNA-binding transcription factor that plays a central role in the differentiation and development of type A pancreatic cells, which are located in the islets of Langerhans and secrete glucagon (By similarity). Binds to specific sequence motif 5'-TAATTA-3' in regulatory elements of target genes (PubMed:22194193, PubMed:31691806). Also plays a key role in intestinal enteroendocrine cell differentiation by promoting differentiation of intestinal enteroendocrine cell types L, -N, -D, -I and -G (PubMed:37883554). Required for normal brain development: plays a role in neuronal proliferation,

	interneuronal migration and differentiation in the embryonic forebrain (PubMed:11889467, PubMed:12379852, PubMed:14722918). May also be involved in axonal guidance in the floor plate (By similarity). Promotes expression of transcription of KDM5C (PubMed:31691806). Activates expression of KDM5C synergistically with histone lysine demethylase PHF8 and perhaps in competition with transcription regulator ZNF711; synergy may be related to enrichment of histone H3K4me3 in regulatory elements (PubMed:31691806).
Cellular Location	Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000255|PROSITE-ProRule:PRU00138, ECO:0000269|PubMed:22194193} Chromosome
Tissue Location	Expressed predominantly in fetal and adult brain and skeletal muscle. Expression is specific to the telencephalon and ventral thalamus. There is an absence of expression in the cerebellum throughout development and also in adult.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.