

## Goat anti-SEPT6, Biotinylated Antibody

Peptide-affinity purified goat antibody Catalog # AF4478a

## **Product Information**

Application	WB, Pep-ELISA
Primary Accession	<u>Q14141</u>
Other Accession	<u>NP_665798.1, NP_055944.2, NP_665801.1</u>
Reactivity	Human, Dog, Bovine
Host	Goat
Clonality	Polyclonal
Clone Names	SEPTIN6
Calculated MW	49717

## **Additional Information**

Gene ID	23157
Other Names	SEPT6; septin 6; KIAA0128; MGC16619; MGC20339; SEP2; SEPT2; septin 2
Dilution	WB~~1:1000 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Immunogen	This antibody is expected to recognize the reported isoforms NP_665798.1, NP_055944.2 and NP_665801.1. Reported variants NP_665799.1 and NP_665798.1 represent identical protein.
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-SEPT6, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **Protein Information**

Name	SEPTIN6 ( <u>HGNC:15848</u> )
Synonyms	KIAA0128, SEP2, SEPT6
Function	Filament-forming cytoskeletal GTPase. Required for normal organization of the actin cytoskeleton. Involved in cytokinesis. May play a role in HCV RNA replication. Forms a filamentous structure with SEPTIN12, SEPTIN6, SEPTIN2 and probably SEPTIN4 at the sperm annulus which is required for the

	structural integrity and motility of the sperm tail during postmeiotic differentiation (PubMed: <u>25588830</u> ).
Cellular Location	Cytoplasm. Cytoplasm, cytoskeleton, spindle Chromosome, centromere, kinetochore Cleavage furrow. Midbody. Cell projection, cilium, flagellum. Note=In metaphase cells, localized within the microtubule spindle. At the metaphase plate, in close apposition to the kinetochores of the congressed chromosomes. In cells undergoing cytokinesis, localized to the midbody, the ingressing cleavage furrow, and the central spindle. Found in the sperm annulus (PubMed:25588830).
Tissue Location	Widely expressed

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.