

Goat anti-SEPT6, Biotinylated Antibody

Peptide-affinity purified goat antibody Catalog # AF4478a

Product Information

Application WB, Pep-ELISA

Primary Accession Q14141

Other Accession NP 665798.1, NP 055944.2, NP 665801.1

Reactivity Human, Dog, Bovine

Host Goat Clonality Polyclonal **Clone Names** SEPTIN6 Calculated MW 49717

Additional Information

Gene ID 23157

Other Names SEPT6; septin 6; KIAA0128; MGC16619; MGC20339; SEP2; SEPT2; septin 2

Dilution WB~~1:1000 Pep-ELISA~~N/A

Format Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5%

bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and

thawing.

This antibody is expected to recognize the reported isoforms NP_665798.1, **Immunogen**

NP_055944.2 and NP_665801.1. Reported variants NP_665799.1 and

NP_665798.1 represent identical protein.

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store Storage

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Goat anti-SEPT6, Biotinylated Antibody is for research use only and not for

use in diagnostic or therapeutic procedures.

Protein Information

Name SEPTIN6 (<u>HGNC:15848</u>)

Synonyms KIAA0128, SEP2, SEPT6

Function Filament-forming cytoskeletal GTPase. Required for normal organization of

> the actin cytoskeleton. Involved in cytokinesis. May play a role in HCV RNA replication. Forms a filamentous structure with SEPTIN12, SEPTIN6, SEPTIN2 and probably SEPTIN4 at the sperm annulus which is required for the

structural integrity and motility of the sperm tail during postmeiotic differentiation (PubMed: <u>25588830</u>).

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton, spindle Chromosome, centromere, kinetochore Cleavage furrow. Midbody. Cell projection, cilium, flagellum. Note=In metaphase cells, localized within the microtubule spindle. At the metaphase plate, in close apposition to the kinetochores of the congressed chromosomes. In cells undergoing cytokinesis, localized to the midbody, the ingressing cleavage furrow, and the central spindle. Found in the sperm annulus (PubMed:25588830).

Tissue Location

Widely expressed..

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.