

Goat anti-HSPC117 (aa201-215), Biotinylated Antibody

Peptide-affinity purified goat antibody Catalog # AF4482a

Product Information

Application WB, Pep-ELISA

Primary Accession Q9Y3I0
Other Accession NP_055121.1

Reactivity Human, Mouse, Rat, Pig, Dog, Bovine

HostGoatClonalityPolyclonalClone NamesC22orf28Calculated MW55210

Additional Information

Gene ID 51493

Other Names C22orf28; chromosome 22 open reading frame 28; DJ149A16.6;

RP1-149A16.6; hypothetical protein LOC51493; novel protein HSPC117

Dilution WB~~1:1000 Pep-ELISA~~N/A

Format Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5%

bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and

thawing.

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Goat anti-HSPC117 (aa201-215), Biotinylated Antibody is for research use only

and not for use in diagnostic or therapeutic procedures.

Protein Information

Name RTCB {ECO:0000255 | HAMAP-Rule:MF_03144}

Synonyms C22orf28

Function Catalytic subunit of the tRNA-splicing ligase complex that acts by directly

joining spliced tRNA halves to mature-sized tRNAs by incorporating the precursor-derived splice junction phosphate into the mature tRNA as a canonical 3',5'-phosphodiester. May act as an RNA ligase with broad substrate

specificity, and may function toward other RNAs.

Nucleus. Cytoplasm {ECO:0000255 | HAMAP-Rule:MF 03144,

Cellular Location

ECO:0000269 | PubMed:24608264} Note=Enters into the nucleus in case of active transcription while it accumulates in cytosol when transcription level is low

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.