

Goat anti-FOXP3 Antibody

Peptide-affinity purified goat antibody

Catalog # AF4493a

Product Information

Application	FC, Pep-ELISA
Primary Accession	Q9BZS1
Other Accession	NP_054728 , NP_001107849.1
Reactivity	Human, Bovine
Host	Goat
Clonality	Polyclonal
Clone Names	FOXP3
Calculated MW	47244

Additional Information

Gene ID	50943
Other Names	FOXP3; forkhead box P3; JM2; AIID; IPEX; PIDX; XPID; DIETER; SCURFIN; scurfin; JM2 protein; immunodeficiency, polyendocrinopathy, enteropathy, X-linked; immune dysregulation, polyendocrinopathy, enteropathy, X-linked
Dilution	FC~~1:10~50 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-FOXP3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FOXP3
Synonyms	IPEX
Function	Transcriptional regulator which is crucial for the development and inhibitory function of regulatory T-cells (Treg) (PubMed: 17377532 , PubMed: 21458306 , PubMed: 23947341 , PubMed: 24354325 , PubMed: 24722479 , PubMed: 24835996 , PubMed: 30513302 , PubMed: 32644293). Plays an essential role in maintaining homeostasis of the immune system by allowing the acquisition of full suppressive function and stability of the Treg lineage, and

by directly modulating the expansion and function of conventional T-cells (PubMed:[23169781](#)). Can act either as a transcriptional repressor or a transcriptional activator depending on its interactions with other transcription factors, histone acetylases and deacetylases (PubMed:[17377532](#), PubMed:[21458306](#), PubMed:[23947341](#), PubMed:[24354325](#), PubMed:[24722479](#)). The suppressive activity of Treg involves the coordinate activation of many genes, including CTLA4 and TNFRSF18 by FOXP3 along with repression of genes encoding cytokines such as interleukin-2 (IL2) and interferon-gamma (IFNG) (PubMed:[17377532](#), PubMed:[21458306](#), PubMed:[23947341](#), PubMed:[24354325](#), PubMed:[24722479](#)). Inhibits cytokine production and T-cell effector function by repressing the activity of two key transcription factors, RELA and NFATC2 (PubMed:[15790681](#)). Mediates transcriptional repression of IL2 via its association with histone acetylase KAT5 and histone deacetylase HDAC7 (PubMed:[17360565](#)). Can activate the expression of TNFRSF18, IL2RA and CTLA4 and repress the expression of IL2 and IFNG via its association with transcription factor RUNX1 (PubMed:[17377532](#)). Inhibits the differentiation of IL17 producing helper T-cells (Th17) by antagonizing RORC function, leading to down-regulation of IL17 expression, favoring Treg development (PubMed:[18368049](#)). Inhibits the transcriptional activator activity of RORA (PubMed:[18354202](#)). Can repress the expression of IL2 and IFNG via its association with transcription factor IKZF4 (By similarity).

Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00089, ECO:0000269 | PubMed:17360565, ECO:0000269 | PubMed:18354202, ECO:0000269 | PubMed:22678915, ECO:0000269 | PubMed:23396208, ECO:0000269 | PubMed:23973222, ECO:0000269 | PubMed:23973223, ECO:0000269 | PubMed:32644293}. Cytoplasm Note=Predominantly expressed in the cytoplasm in activated conventional T-cells whereas predominantly expressed in the nucleus in regulatory T- cells (Treg). The 41 kDa form derived by proteolytic processing is found exclusively in the chromatin fraction of activated Treg cells (By similarity). {ECO:0000250 | UniProtKB:Q99JB6, ECO:0000269 | PubMed:22678915}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.