

# Goat anti-SOX2 Antibody

Peptide-affinity purified goat antibody Catalog # AF4507a

#### **Product Information**

**Application** WB, IF, FC, Pep-ELISA

Primary Accession P48431
Other Accession NP\_003097.1

**Reactivity** Human, Mouse, Rat, Dog, Bovine

HostGoatClonalityPolyclonalClone NamesSOX2Calculated MW34310

### **Additional Information**

**Gene ID** 6657

Other Names SOX2; SRY (sex determining region Y)-box 2; ANOP3; MGC2413; SRY-related

HMG-box gene 2; sex-determining region Y-box 2; transcription factor SOX2

**Dilution** WB~~1:1000 IF~~1:50~200 FC~~1:10~50 Pep-ELISA~~N/A

**Format** Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5%

bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and

thawing.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** Goat anti-SOX2 Antibody is for research use only and not for use in diagnostic

or therapeutic procedures.

## **Protein Information**

Name SOX2

**Function** Transcription factor that forms a trimeric complex with OCT4 on DNA and

controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206 (By similarity). Binds to the proximal enhancer region of NANOG (By similarity). Critical for early

embryogenesis and for embryonic stem cell pluripotency

(PubMed: 18035408). Downstream SRRT target that mediates the promotion of

neural stem cell self-renewal (By similarity). Keeps neural cells

undifferentiated by counteracting the activity of proneural proteins and suppresses neuronal differentiation (By similarity). May function as a switch in

neuronal development (By similarity).

### **Cellular Location**

Nucleus speckle {ECO:0000250|UniProtKB:Q05066}. Cytoplasm

{ECO:0000250|UniProtKB:Q05738}. Nucleus

{ECO:0000250 | UniProtKB:Q05738}. Note=Acetylation contributes to its nuclear localization and deacetylation by HDAC3 induces a cytoplasmic delocalization (By similarity). Colocalizes in the nucleus with ZNF208 isoform KRAB-O and tyrosine hydroxylase (TH) (By similarity) Colocalizes with SOX6 in speckles. Colocalizes with CAML in the nucleus (By similarity). Nuclear import is facilitated by XPO4, a protein that usually acts as a nuclear export signal

receptor (By similarity) {ECO:0000250 | UniProtKB:Q05066,

ECO:0000250 | UniProtKB:Q05738}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.