

# Goat anti-VE-cadherin Antibody

Peptide-affinity purified goat antibody

Catalog # AF4510a

## Product Information

---

<b>Application</b>	IF, FC, Pep-ELISA
<b>Primary Accession</b>	<a href="#">P33151</a>
<b>Other Accession</b>	<a href="#">NP_001786.2</a>
<b>Reactivity</b>	Human, Pig, Bovine
<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal
<b>Clone Names</b>	CDH5
<b>Calculated MW</b>	87528

## Additional Information

---

<b>Gene ID</b>	1003
<b>Other Names</b>	CDH5; cadherin 5, type 2 (vascular endothelium); 7B4; CD144; 7B4 antigen; VE-cadherin; cadherin 5, type 2, VE-cadherin (vascular epithelium); cadherin-5; cd144 antigen; endothelial-specific cadherin; vascular endothelial cadherin
<b>Dilution</b>	IF~~1:50~200 FC~~1:10~50 Pep-ELISA~~N/A
<b>Format</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Goat anti-VE-cadherin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CDH5 ( <a href="#">HGNC:1764</a> )
<b>Function</b>	Cadherins are calcium-dependent cell adhesion proteins (By similarity). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types (PubMed: <a href="#">21269602</a> ). This cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions (By similarity). It associates with alpha-catenin forming a link to the cytoskeleton (PubMed: <a href="#">10861224</a> ). Plays a role in coupling actin fibers to cell junctions in endothelial cells, via acting as a cell

junctional complex anchor for AMOTL2 and MAGI1 (By similarity). Acts in concert with KRIT1 and PALS1 to establish and maintain correct endothelial cell polarity and vascular lumen (By similarity). These effects are mediated by recruitment and activation of the Par polarity complex and RAP1B (PubMed:[20332120](#)). Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction (PubMed:[20332120](#)). Associates with CTNND1/p120-catenin to control CADH5 endocytosis (By similarity).

**Cellular Location**

Cell junction, adherens junction. Cell membrane; Single-pass type I membrane protein. Cytoplasm {ECO:0000250|UniProtKB:P55284}. Note=Found at cell-cell boundaries and probably at cell-matrix boundaries. KRIT1 and CDH5 reciprocally regulate their localization to endothelial cell-cell junctions

**Tissue Location**

Expressed in endothelial cells (at protein level) (PubMed:27338829). Expressed in the brain (PubMed:2059658)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.