

Goat anti-OCIL Antibody

Peptide-affinity purified goat antibody Catalog # AF4513a

Product Information

Application WB, IHC, IF, FC, Pep-ELISA

Primary Accession Q9UHP7

Other Accession NP 037401.1, NP 001004419.1, NP 001184246.1

Reactivity Human, Dog, Bovine

HostGoatClonalityPolyclonalClone NamesCLEC2DCalculated MW21849

Additional Information

Gene ID 29121

Other Names CLEC2D; C-type lectin domain family 2, member D; CLAX; LLT1; OCIL; C-type

lectin related f; C-type lectin superfamily 2, member D; lectin-like NK cell

receptor; lectin-like transcript 1; osteoclast inhibitory lectin

Dilution WB~~1:1000 IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50 Pep-ELISA~~N/A

Format Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5%

bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and

thawing.

Immunogen This antibody is expected to recognize all isoforms (NP_037401.1;

NP 001004419.1; NP 001184246.1).

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Goat anti-OCIL Antibody is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name CLEC2D

Synonyms CLAX, LLT1, OCIL

Function Receptor for KLRB1 that protects target cells against natural killer

cell-mediated lysis (PubMed:<u>16339513</u>, PubMed:<u>20843815</u>). Inhibits

osteoclast formation (PubMed: 14753741, PubMed: 15123656). Inhibits bone

resorption (PubMed: 14753741). Modulates the release of interferon-gamma

(PubMed:15104121). Binds high molecular weight sulfated

glycosaminoglycans (PubMed: 15123656).

Cellular Location Cell membrane; Single-pass type II membrane protein [Isoform 4]:

Endoplasmic reticulum

Tissue Location Detected in peripheral blood leukocytes, osteoblasts, lymph node, thymus

and spleen. Isoform 1, isoform 2 and isoform 4 are expressed in T- and B-lymphocytes, and at lower levels in NK cells. They are also expressed in

B-cell lines and LPS-matured monocyte-derived dendritic cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.