

# Goat anti-SCRIB / Scribble Antibody

Peptide-affinity purified goat antibody

Catalog # AF4525a

## Product Information

---

Application	WB, IF, Pep-ELISA
Primary Accession	<a href="#">Q14160</a>
Other Accession	<a href="#">NP_874365.3</a> , <a href="#">NP_056171.3</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Clone Names	SCRIB
Calculated MW	177724

## Additional Information

---

Gene ID	23513
Other Names	SCRIB; scribbled homolog (Drosophila); CRIB1; SCRIB1; SCRIB1; Vartul; scribble
Dilution	WB~~1:1000 IF~~1:50~200 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Immunogen	This antibody is expected to recognize both reported isoforms (NP_874365.3; NP_056171.3).
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-SCRIB / Scribble Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	SCRIB ( <a href="#">HGNC:30377</a> )
Function	Scaffold protein involved in different aspects of polarized cell differentiation regulating epithelial and neuronal morphogenesis and T-cell polarization (PubMed: <a href="#">15182672</a> , PubMed: <a href="#">16344308</a> , PubMed: <a href="#">16965391</a> , PubMed: <a href="#">18641685</a> , PubMed: <a href="#">18716323</a> , PubMed: <a href="#">19041750</a> , PubMed: <a href="#">27380321</a> ). Via its interaction with CRTAM, required for the late phase polarization of a subset of CD4+ T-cells, which in turn regulates TCR-mediated proliferation and IFNG and IL22 production (By similarity).

Plays a role in cell directional movement, cell orientation, cell sheet organization and Golgi complex polarization at the cell migration front (By similarity). Promotes epithelial cell layer barrier function via maintaining cell-cell adhesion (By similarity). Most probably functions in the establishment of apico- basal cell polarity (PubMed:[16344308](#), PubMed:[19041750](#)). May function in cell proliferation regulating progression from G1 to S phase and as a positive regulator of apoptosis for instance during acinar morphogenesis of the mammary epithelium (PubMed:[16965391](#), PubMed:[19041750](#)). May regulate cell invasion via MAPK-mediated cell migration and adhesion (PubMed:[18641685](#), PubMed:[18716323](#)). May play a role in exocytosis and in the targeting of synaptic vesicles to synapses (PubMed:[15182672](#)). Functions as an activator of Rac GTPase activity (PubMed:[15182672](#)).

#### Cellular Location

Cell membrane; Peripheral membrane protein. Cell junction. Cell junction, adherens junction. Cell projection, lamellipodium. Cytoplasm. Postsynapse. Presynapse. Note=Targeting to cell-cell junctions which is CDH1-dependent is required for the pro-apoptotic activity. In a subset of CD4+ T-cells, colocalizes with CRTAM at the immunological synapse during the late phase of T-cell activation (By similarity) Localized to small puncta throughout the cytoplasm and cell membrane when in the presence of SNAIL1 (By similarity). Localized along the length of perinuclear emanating vimentin bundles and at vimentin-positive fibrils at the cell periphery (PubMed:19386766). Localized to the lateral plasma membrane during the establishment and maturation of cell-cell contacts (PubMed:19386766) {ECO:0000250|UniProtKB:A0A8P0N4K0, ECO:0000250|UniProtKB:Q80U72, ECO:0000269|PubMed:19386766}

#### Tissue Location

Expressed in kidney, skeletal muscles, liver, lung, breast, intestine, placenta and skin mainly in epithelial cells (at protein level).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.