

Goat anti-Ku70 / G22P1 Antibody

Peptide-affinity purified goat antibody

Catalog # AF4528a

Product Information

Application	FC, Pep-ELISA
Primary Accession	P12956
Other Accession	NP_001460.1 , NP_001275907.1
Reactivity	Human, Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Clone Names	XRCC6
Calculated MW	69843

Additional Information

Gene ID	2547
Other Names	G22P1; thyroid autoantigen 70kDa (Ku antigen); HGNC:4055; CTC75; CTCBF; KU70; ML8; TLAA; ATP-dependent DNA helicase II, 70 kDa subunit; CTC box binding factor 75 kDa subunit; OTTHUMP00000028581; thyroid autoantigen 70kD (Ku antigen); thyroid-lupus autoant
Dilution	FC~1:10~50 Pep-ELISA~~N/A
Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Goat anti-Ku70 / G22P1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	XRCC6
Synonyms	G22P1
Function	Single-stranded DNA-dependent ATP-dependent helicase that plays a key role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed: 11493912 , PubMed: 12145306 , PubMed: 20493174 , PubMed: 2466842 , PubMed: 7957065 , PubMed: 8621488 , PubMed: 9742108). Required for double-strand break repair and V(D) recombination

(PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#),
PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)).
Also has a role in chromosome translocation (PubMed:[11493912](#),
PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#),
PubMed:[8621488](#), PubMed:[9742108](#)). Has a role in chromosome translocation
(PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#),
PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)).
The DNA helicase II complex binds preferentially to fork-like ends of
double-stranded DNA in a cell cycle-dependent manner (PubMed:[11493912](#),
PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#),
PubMed:[8621488](#), PubMed:[9742108](#)). It works in the 3'-5' direction
(PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#),
PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)).
During NHEJ, the XRCC5-XRRC6 dimer performs the recognition step: it
recognizes and binds to the broken ends of the DNA and protects them from
further resection (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#),
PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)).
Binding to DNA may be mediated by XRCC6 (PubMed:[11493912](#),
PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#),
PubMed:[8621488](#), PubMed:[9742108](#)). The XRCC5-XRRC6 dimer acts as a
regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by
increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold
(PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#),
PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)).
The XRCC5-XRRC6 dimer is probably involved in stabilizing broken DNA ends
and bringing them together (PubMed:[11493912](#), PubMed:[12145306](#),
PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#),
PubMed:[9742108](#)). The assembly of the DNA-PK complex to DNA ends is
required for the NHEJ ligation step (PubMed:[11493912](#), PubMed:[12145306](#),
PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#),
PubMed:[9742108](#)). Probably also acts as a 5'-deoxyribose-5-phosphate lyase
(5'-dRP lyase), by catalyzing the beta-elimination of the 5'
deoxyribose-5-phosphate at an abasic site near double-strand breaks
(PubMed:[20383123](#)). 5'-dRP lyase activity allows to 'clean' the termini of
abasic sites, a class of nucleotide damage commonly associated with strand
breaks, before such broken ends can be joined (PubMed:[20383123](#)). The
XRCC5-XRRC6 dimer together with APEX1 acts as a negative regulator of
transcription (PubMed:[8621488](#)). In association with NAA15, the XRCC5-XRRC6
dimer binds to the osteocalcin promoter and activates osteocalcin expression
(PubMed:[12145306](#)). Plays a role in the regulation of DNA virus-mediated
innate immune response by assembling into the HDP-RNP complex, a
complex that serves as a platform for IRF3 phosphorylation and subsequent
innate immune response activation through the cGAS-STING pathway
(PubMed:[28712728](#)). Negatively regulates apoptosis by interacting with BAX
and sequestering it from the mitochondria (PubMed:[15023334](#)). Might have
deubiquitination activity, acting on BAX (PubMed:[18362350](#)).

Cellular Location

Nucleus. Chromosome. Cytoplasm. Note=When trimethylated, localizes in the cytoplasm.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.