

# Goat anti-TLR4 / CD284 Antibody

Peptide-affinity purified goat antibody  
Catalog # AF4536a

## Product Information

---

<b>Application</b>	IHC, IF, FC, Pep-ELISA
<b>Primary Accession</b>	<a href="#">O00206</a>
<b>Other Accession</b>	<a href="#">NP_612564.1</a>
<b>Reactivity</b>	Human, Mouse, Rat, Pig, Dog, Sheep, Bovine
<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal
<b>Clone Names</b>	TLR4

## Additional Information

---

<b>Other Names</b>	TLR4; toll-like receptor 4; ARMD10; CD284; TOLL; hToll; OTTHUMP00000022807; homolog of Drosophila toll
<b>Dilution</b>	IHC~~1:100~500 IF~~1:50~200 FC~~1:10~50 Pep-ELISA~~N/A
<b>Format</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Goat anti-TLR4 / CD284 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



## Protein Information

---

## Images

---

EB09441 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing Golgi/ER and cytoplasmic staining. The nuclear s



EB09441 Flow cytometric analysis of paraformaldehyde fixed U937 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) fol

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.