

KD-Validated Anti-KEAP1 Mouse Monoclonal Antibody

Mouse monoclonal antibody

Catalog # AGI1111

Product Information

Application	WB, FC
Primary Accession	Q14145
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Clone Names	25GB495
Calculated MW	69666
Gene Name	KEAP1
Aliases	KEAP1; Kelch Like ECH Associated Protein 1; KLHL19; INrf2; KIAA0132; Kelch-Like ECH-Associated Protein 1; Kelch-Like Family Member 19; Cytosolic Inhibitor Of Nrf2; Kelch-Like Protein 19; MGC10630; MGC20887; MGC1114; MGC4407; MGC9454; KEAP1 Delta C; INRF2
Immunogen	Recombinant protein of human KEAP1

Additional Information

Gene ID	9817
Other Names	Kelch-like ECH-associated protein 1, Cytosolic inhibitor of Nrf2 {ECO:0000303 Ref.1}, INrf2 {ECO:0000303 Ref.1}, Kelch-like protein 19 {ECO:0000312 HGNC:HGNC:23177}, KEAP1 {ECO:0000303 PubMed:14585973, ECO:0000312 HGNC:HGNC:23177}

Protein Information

Name	KEAP1 {ECO:0000303 PubMed:14585973, ECO:0000312 HGNC:HGNC:23177}
Function	Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin ligase complex that regulates the response to oxidative stress by targeting NFE2L2/NRF2 for ubiquitination (PubMed: 14585973 , PubMed: 15379550 , PubMed: 15572695 , PubMed: 15601839 , PubMed: 15983046 , PubMed: 37339955). KEAP1 acts as a key sensor of oxidative and electrophilic stress: in normal conditions, the BCR(KEAP1) complex mediates ubiquitination and degradation of NFE2L2/NRF2, a transcription factor regulating expression of many cytoprotective genes (PubMed: 15601839 , PubMed: 16006525). In response to oxidative stress, different electrophile metabolites trigger non-enzymatic covalent modifications of highly reactive cysteine residues in KEAP1, leading to inactivate the ubiquitin ligase activity of the BCR(KEAP1) complex, promoting NFE2L2/NRF2 nuclear accumulation and expression of phase II detoxifying enzymes (PubMed: 16006525 , PubMed: 17127771 , PubMed: 18251510 , PubMed: 19489739 , PubMed: 29590092). In response to

selective autophagy, KEAP1 is sequestered in inclusion bodies following its interaction with SQSTM1/p62, leading to inactivation of the BCR(KEAP1) complex and activation of NFE2L2/NRF2 (PubMed:[20452972](#)). The BCR(KEAP1) complex also mediates ubiquitination of SQSTM1/p62, increasing SQSTM1/p62 sequestering activity and degradation (PubMed:[28380357](#)). The BCR(KEAP1) complex also targets BPTF and PGAM5 for ubiquitination and degradation by the proteasome (PubMed:[15379550](#), PubMed:[17046835](#)).

Cellular Location

Cytoplasm. Nucleus. Note=Mainly cytoplasmic (PubMed:15601839). In response to selective autophagy, relocalizes to inclusion bodies following interaction with SQSTM1/p62 (PubMed:20452972).

Tissue Location

Broadly expressed, with highest levels in skeletal muscle.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.