

KD-Validated Anti-CD13 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1179

Product Information

Application	WB, FC, ICC
Primary Accession	P15144
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB3360
Calculated MW	109540
Gene Name	ANPEP
Aliases	ANPEP; Alanyl Aminopeptidase, Membrane; AP-N; HAPN; Microsomal Aminopeptidase; Aminopeptidase N; Aminopeptidase M; LAP1; P150; CD13; PEPN; Myeloid Plasma Membrane Glycoprotein CD13; Alanyl (Membrane) Aminopeptidase; Membrane Alanyl Aminopeptidase; EC 3.4.11.2; Gp150; GP150; AP-M; APN; Alanyl; Aminopeptidase; CD13 Antigen; EC 3.4.11
Immunogen	A synthesized peptide derived from human CD13

Additional Information

Gene ID	290
Other Names	Aminopeptidase N, AP-N, hAPN, 3.4.11.2, Alanyl aminopeptidase, Aminopeptidase M, AP-M, Microsomal aminopeptidase, Myeloid plasma membrane glycoprotein CD13, gp150, CD13, ANPEP, APN, CD13, PEPN

Protein Information

Name	ANPEP
Synonyms	APN, CD13, PEPN
Function	Broad specificity aminopeptidase which plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Also involved in the processing of various peptides including peptide hormones, such as angiotensin III and IV, neuropeptides, and chemokines. May also be involved the cleavage of peptides bound to major histocompatibility complex class II molecules of antigen presenting cells. May have a role in angiogenesis and promote cholesterol crystallization. May have a role in amino acid transport by acting as binding partner of amino acid transporter SLC6A19 and regulating its activity (By similarity).
Cellular Location	Cell membrane; Single-pass type II membrane protein. Note=Also found as a soluble form

Tissue Location

Expressed in epithelial cells of the kidney, intestine, and respiratory tract; granulocytes, monocytes, fibroblasts, endothelial cells, cerebral pericytes at the blood-brain barrier, synaptic membranes of cells in the CNS. Also expressed in endometrial stromal cells, but not in the endometrial glandular cells. Found in the vasculature of tissues that undergo angiogenesis and in malignant gliomas and lymph node metastases from multiple tumor types but not in blood vessels of normal tissues. A soluble form has been found in plasma. It is found to be elevated in plasma and effusions of cancer patients.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.