

KD-Validated Anti-CDC73 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1207

Product Information

Application	WB, FC, ICC
Primary Accession	Q6P1J9
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB3450
Calculated MW	60577
Gene Name	CDC73
Aliases	CDC73; Cell Division Cycle 73; Parafibromin; HRPT2; FIHP; Paf1/RNA Polymerase II Complex Component; Cell Division Cycle Protein 73 Homolog; Familial Isolated Hyperparathyroidism; Hyperparathyroidism 2 Protein; C1orf28; HRPT1; Cell Division Cycle 73, Paf1/RNA Polymerase II Complex Component, Homolog (S. Cerevisiae); Cell Division Cycle 73 Paf1/RNA Polymerase II Complex Component-Like Protein; Cell Division Cycle 73, Paf1/RNA Polymerase II Complex Component, Homolog; Hyperparathyroidism 2 (With Jaw Tumor); Chromosome 1 Open Reading Frame 28; Hyperparathyroidism 1; PARAFIBROMIN; C1ORF28; HPTJT; HYX
Immunogen	A synthesized peptide derived from human HRPT2 / CDC73

Additional Information

Gene ID	79577
Other Names	Parafibromin, Cell division cycle protein 73 homolog, Hyperparathyroidism 2 protein, CDC73, C1orf28, HRPT2

Protein Information

Name	CDC73
Synonyms	C1orf28, HRPT2
Function	Tumor suppressor probably involved in transcriptional and post-transcriptional control pathways. May be involved in cell cycle progression through the regulation of cyclin D1/PRAD1 expression. Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser- 5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and

HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1; it promotes leukemogenesis through association with KMT2A/MLL1-rearranged oncoproteins, such as KMT2A/MLL1-MLLT3/AF9 and KMT2A/MLL1-MLLT1/ENL. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1); UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. In case of infection by influenza A strain H3N2, PAF1C associates with viral NS1 protein, thereby regulating gene transcription. Connects PAF1C with the cleavage and polyadenylation specificity factor (CPSF) complex and the cleavage stimulation factor (CSTF) complex, and with Wnt signaling. Involved in polyadenylation of mRNA precursors.

Cellular Location

Nucleus

Tissue Location

Found in adrenal and parathyroid glands, kidney and heart.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.