

KD-Validated Anti-Histone deacetylase 8 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody
Catalog # AGI1211

Product Information

Application	WB, FC, ICC
Primary Accession	Q9BY41
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	25GB2495
Calculated MW	41758
Gene Name	HDAC8
Aliases	HDAC8; Histone Deacetylase 8; HDACL1; KDAC8; RPD3; Protein Decrotonylase HDAC8; Histone Deacetylase-Like 1; Protein Deacetylase HDAC8; MRXS6; HD8; WTS; Wilson-Turner X-Linked Mental Retardation Syndrome; EC 3.5.1.98; EC 3.5.1.-; CDA07; CDLS5
Immunogen	A synthesized peptide derived from human PKM2

Additional Information

Gene ID	55869
Other Names	Histone deacetylase 8, HD8, 3.5.1.98, Protein deacetylase HDAC8, 3.5.1.-, Protein decrotonylase HDAC8, 3.5.1.-, HDAC8 {ECO:0000303 PubMed:10926844, ECO:0000312 HGNC:HGNC:13315}

Protein Information

Name	HDAC8 {ECO:0000303 PubMed:10926844, ECO:0000312 HGNC:HGNC:13315}
Function	Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed: 10748112 , PubMed: 10922473 , PubMed: 10926844 , PubMed: 14701748 , PubMed: 28497810). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed: 10748112 , PubMed: 10922473 , PubMed: 10926844 , PubMed: 14701748). Histone deacetylases act via the formation of large multiprotein complexes (PubMed: 10748112 , PubMed: 10922473 , PubMed: 10926844 , PubMed: 14701748). Also involved in the deacetylation of cohesin complex protein SMC3 regulating release of cohesin complexes from chromatin (PubMed: 22885700). May play a role in smooth muscle cell contractility (PubMed: 15772115). In addition to protein deacetylase activity,

also has protein-lysine deacylase activity: acts as a protein decrotonylase by mediating decrotonylation ((2E)-butenoyl) of histones (PubMed:[28497810](#)).

Cellular Location

Nucleus. Chromosome Cytoplasm Note=Excluded from the nucleoli (PubMed:10748112). Found in the cytoplasm of cells showing smooth muscle differentiation (PubMed:15772115, PubMed:16538051).

Tissue Location

Weakly expressed in most tissues. Expressed at higher level in heart, brain, kidney and pancreas and also in liver, lung, placenta, prostate and kidney.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.