

KD-Validated Anti-ACADVL Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1225

Product Information

Application	WB, FC, ICC
Primary Accession	P49748
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB3615
Calculated MW	70390
Gene Name	ACADVL
Aliases	ACADVL; Acyl-CoA Dehydrogenase Very Long Chain; VLCAD; LCACD; ACAD6; Very Long-Chain Specific Acyl-CoA Dehydrogenase, Mitochondrial; Acyl-Coenzyme A Dehydrogenase, Very Long Chain; EC 1.3.8.9; EC 1.3.99
Immunogen	A synthesized peptide derived from human ACADVL/VLCAD

Additional Information

Gene ID	37
Other Names	Very long-chain specific acyl-CoA dehydrogenase, mitochondrial, VLCAD, 1.3.8.9, ACADVL (HGNC:92)

Protein Information

Name	ACADVL (HGNC:92)
Function	Catalyzes the first of the four reactions of the mitochondrial fatty acid beta-oxidation (FAO) pathway, which consists in the proR-proR stereospecific alpha, beta-dehydrogenation of fatty acyl-CoA thioesters using the electron transfer flavoprotein (ETF) as their physiologic electron acceptor, resulting in the formation of trans-2-enoyl-CoA ((2E)-enoyl-CoA) (PubMed:17564966, PubMed:18227065, PubMed:26474213, PubMed:7668252, PubMed:9461620, PubMed:9599005, PubMed:9839948). The mitochondrial FAO pathway is the major energy- producing process in tissues and is performed through cycles of four consecutive reactions (PubMed:26474213, PubMed:7668252). Each FAO cycle shortens the fatty acyl-CoA by two carbons, yielding one acetyl-CoA (for the citric acid cycle), one FADH(2), and one NADH (which donate electrons to the respiratory chain for ATP production) (PubMed:26474213, PubMed:7668252). Among the different mitochondrial acyl-CoA dehydrogenases, very long-chain specific acyl-CoA dehydrogenase acts specifically on fatty acyl-CoAs with saturated 12 to 24 carbons long primary chains (PubMed:17564966, PubMed:21237683, PubMed:9839948), but can also catalyze monounsaturated fatty acids such as oleate ((9Z)-octadecenoate), (9Z)-hexadecenoate, and others (PubMed:17564966).

Can use (4Z,7Z,10Z,13Z,16Z,19Z)-docosahexaenoate as substrate in vitro (which is not primarily used for energy but mainly beta-oxidized in the peroxisomes) (PubMed:[17564966](#), PubMed:[26474213](#)). In addition, based on its established catalytic mechanism, and combined genetic interaction or mutant phenotype evidence, it is predicted to act also on substrates that have not been tested experimentally but are metabolized by mitochondrial FAO, including long-chain unsaturated fatty acids such as linoleate (9Z,12Z-octadecadienoate), linolenate (9Z,12Z,15Z-octadecatrienoate), and others (PubMed:[26474213](#)). Among the different mitochondrial acyl-CoA dehydrogenases, its FAO activity overlaps with that of ACAD9 and ACADL, but plays a primary role in tissues where it is the main long-chain ACAD expressed, such as the heart and skeletal muscle (PubMed:[17564966](#)).

Cellular Location

Mitochondrion inner membrane; Peripheral membrane protein

Tissue Location

Predominantly expressed in heart and skeletal muscle (at both mRNA and protein levels) (PubMed:[17564966](#), PubMed:[8845838](#)). Also detected in kidney and liver (at protein level) (PubMed:[8845838](#)).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.