

KD-Validated Anti-TAR DNA binding protein Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1287

Product Information

Application	WB, FC, ICC
Primary Accession	Q13148
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB 2405
Calculated MW	44740
Gene Name	TARDBP
Aliases	TARDBP; TAR DNA Binding Protein; TDP-43; ALS10; TAR DNA-Binding Protein 43; TAR DNA-Binding Protein-43; TDP43
Immunogen	A synthesized peptide derived from human TDP43

Additional Information

Gene ID	23435
Other Names	TAR DNA-binding protein 43, TDP-43, TARDBP {ECO:0000303 PubMed:18396105, ECO:0000312 HGNC:HGNC:11571}

Protein Information

Name	TARDBP {ECO:0000303 PubMed:18396105, ECO:0000312 HGNC:HGNC:11571}
Function	RNA-binding protein that is involved in various steps of RNA biogenesis and processing (PubMed: 23519609). Preferentially binds, via its two RNA recognition motifs RRM1 and RRM2, to GU-repeats on RNA molecules predominantly localized within long introns and in the 3'UTR of mRNAs (PubMed: 23519609 , PubMed: 24240615 , PubMed: 24464995). In turn, regulates the splicing of many non-coding and protein-coding RNAs including proteins involved in neuronal survival, as well as mRNAs that encode proteins relevant for neurodegenerative diseases (PubMed: 21358640 , PubMed: 29438978). Plays a role in maintaining mitochondrial homeostasis by regulating the processing of mitochondrial transcripts (PubMed: 28794432). Also regulates mRNA stability by recruiting CNOT7/CAF1 deadenylase on mRNA 3'UTR leading to poly(A) tail deadenylation and thus shortening (PubMed: 30520513). In response to oxidative insult, associates with stalled ribosomes localized to stress granules (SGs) and contributes to cell survival (PubMed: 19765185 , PubMed: 23398327). Also participates in the normal skeletal muscle formation and regeneration, forming cytoplasmic

myo-granules and binding mRNAs that encode sarcomeric proteins (PubMed:[30464263](#)). Plays a role in the maintenance of the circadian clock periodicity via stabilization of the CRY1 and CRY2 proteins in a FBXL3-dependent manner (PubMed:[27123980](#)). Negatively regulates the expression of CDK6 (PubMed:[19760257](#)). Regulates the expression of HDAC6, ATG7 and VCP in a PPIA/CYPA-dependent manner (PubMed:[25678563](#)).

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, Stress granule Mitochondrion.
Note=Continuously travels in and out of the nucleus (PubMed:18957508).
Localizes to stress granules in response to oxidative stress (PubMed:19765185). A small subset localizes in mitochondria (PubMed:28794432).

Tissue Location

Ubiquitously expressed. In particular, expression is high in pancreas, placenta, lung, genital tract and spleen

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.