

KD-Validated Anti-Serine/Threonine Kinase 39 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1403

Product Information

Application	WB, FC
Primary Accession	Q9UEW8
Reactivity	Rat, Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	25GB3395
Calculated MW	59474
Gene Name	STK39
Aliases	STK39; Serine/Threonine Kinase 39; DCHT; SPAK; STE20/SPS1-Related Proline-Alanine-Rich Protein Kinase; Serine/Threonine-Protein Kinase 39; Ste-20-Related Kinase; EC 2.7.11.1; PASK; Serine Threonine Kinase 39 (STE20/SPS1 Homolog, Yeast); Proline-Alanine-Rich STE20-Related Kinase; Small Intestine SPAK-Like Kinase; STE20/SPS1 Homolog (Yeast); Serine Threonine Kinase 39; Ste20-Like Protein Kinase; STE20/SPS1 Homolog; EC 2.7.11
Immunogen	A synthesized peptide derived from human STK39

Additional Information

Gene ID	27347
Other Names	STE20/SPS1-related proline-alanine-rich protein kinase, Ste-20-related kinase, 2.7.11.1, DCHT, Serine/threonine-protein kinase 39, STK39

Protein Information

Name	STK39
Function	Effector serine/threonine-protein kinase component of the WNK-SPAK/OSR1 kinase cascade, which is involved in various processes, such as ion transport, response to hypertonic stress and blood pressure (PubMed: 16669787 , PubMed: 18270262 , PubMed: 21321328 , PubMed: 34289367). Specifically recognizes and binds proteins with a RFXV motif (PubMed: 16669787 , PubMed: 21321328). Acts downstream of WNK kinases (WNK1, WNK2, WNK3 or WNK4): following activation by WNK kinases, catalyzes phosphorylation of ion cotransporters, such as SLC12A1/NKCC2, SLC12A2/NKCC1, SLC12A3/NCC, SLC12A5/KCC2 or SLC12A6/KCC3, regulating their activity (PubMed: 21321328). Mediates regulatory volume increase in response to hyperosmotic stress by catalyzing phosphorylation of ion cotransporters SLC12A1/NKCC2, SLC12A2/NKCC1 and SLC12A6/KCC3 downstream of WNK1 and WNK3 kinases

(PubMed:[12740379](#), PubMed:[16669787](#), PubMed:[21321328](#)). Phosphorylation of Na-K-Cl cotransporters SLC12A2/NKCC1 and SLC12A2/NKCC1 promote their activation and ion influx; simultaneously, phosphorylation of K-Cl cotransporters SLC12A5/KCC2 and SLC12A6/KCC3 inhibit their activity, blocking ion efflux (PubMed:[16669787](#), PubMed:[19665974](#), PubMed:[21321328](#)). Acts as a regulator of NaCl reabsorption in the distal nephron by mediating phosphorylation and activation of the thiazide-sensitive Na-Cl cotransporter SLC12A3/NCC in distal convoluted tubule cells of kidney downstream of WNK4 (PubMed:[18270262](#)). Mediates the inhibition of SLC4A4, SLC26A6 as well as CFTR activities (By similarity). Phosphorylates RELT (By similarity).

Cellular Location

Cytoplasm. Nucleus. Note=Nucleus when caspase-cleaved.

Tissue Location

Predominantly expressed in brain and pancreas followed by heart, lung, kidney, skeletal muscle, liver, placenta and testis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.