

KD-Validated Anti-Cyclin Dependent Kinase Inhibitor 2A Rabbit Monoclonal Antibody

Rabbit monoclonal antibody
Catalog # AGI1412

Product Information

Application	WB, FC, ICC
Primary Accession	P42771
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	25GB3420
Calculated MW	16533
Gene Name	CDKN2A
Aliases	CDKN2A; Cyclin Dependent Kinase Inhibitor 2A; P14ARF; CDK4I; MTS1; ARF; Cyclin-Dependent Kinase 4 Inhibitor A; P16-INK4A; P16INK4a; P19Arf; INK4a; CDKN2; CMM2; INK4; P16; P19; P14; MLM; Cyclin-Dependent Kinase Inhibitor 2A (Melanoma, P16,Inhibits CDK4); Cyclin-Dependent Kinase Inhibitor 2A; Multiple Tumour Suppressor 1; Multiple Tumor Suppressor 1; CDKN2A/ARF Intron 2 LncRNA; Alternative Reading Frame; Inhibitor Of Cdk4 A; MTS-1; CAI2; Cell Cycle Negative Regulator Beta; P19 Alternate Open Reading Frame; P14 Alternate Open Reading Frame; CDK4 Inhibitor P16-INK4; Tumor Suppressor ARF; P16-INK4a; P16-INK4; P16INK4A; P16INK4; TP16
Immunogen	A synthesized peptide derived from human p16 INK

Additional Information

Gene ID	1029
Other Names	Cyclin-dependent kinase inhibitor 2A {ECO:0000312 HGNC:HGNC:1787}, Cyclin-dependent kinase 4 inhibitor A, CDK4I, Multiple tumor suppressor 1, MTS-1, p16-INK4a, p16-INK4, p16INK4A, CDKN2A (HGNC:1787), CDKN2, MTS1

Protein Information

Name	CDKN2A (HGNC:1787)
Synonyms	CDKN2, MTS1
Function	Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Widely expressed but not detected in brain or skeletal muscle. Isoform 3 is

pancreas-specific

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.