

KD-Validated Anti-AKT3 Mouse Monoclonal Antibody

Mouse monoclonal antibody

Catalog # AGI1441

Product Information

Application	WB, FC
Primary Accession	Q9Y243
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Mouse IgG3
Clone Names	25GB3680
Calculated MW	55775
Gene Name	AKT3
Aliases	AKT3; AKT Serine/Threonine Kinase 3; PKBG; RAC-Gamma; PRKBG; RAC-Gamma Serine/Threonine-Protein Kinase; RAC-PK-Gamma; EC 2.7.11.1; PKB Gamma; STK-2; V-Akt Murine Thymoma Viral Oncogene Homolog 3 (Protein Kinase B, Gamma); V-Akt Murine Thymoma Viral Oncogene Homolog 3; RAC-Gamma Serine/Threonine Protein Kinase; Protein Kinase B, Gamma; Protein Kinase B Gamma; Protein Kinase Akt-3; PKB-GAMMA; EC 2.7.11; MPPH2; MPPH
Immunogen	Recombinant protein of human AKT3

Additional Information

Gene ID	10000
Other Names	RAC-gamma serine/threonine-protein kinase, 2.7.11.1, Protein kinase Akt-3, Protein kinase B gamma, PKB gamma, RAC-PK-gamma, STK-2, AKT3, PKBG

Protein Information

Name	AKT3
Synonyms	PKBG
Function	AKT3 is one of 3 closely related serine/threonine-protein kinases (AKT1, AKT2 and AKT3) called the AKT kinase, and which regulate many processes including metabolism, proliferation, cell survival, growth and angiogenesis. This is mediated through serine and/or threonine phosphorylation of a range of downstream substrates. Over 100 substrate candidates have been reported so far, but for most of them, no isoform specificity has been reported. AKT3 is the least studied AKT isoform. It plays an important role in brain development and is crucial for the viability of malignant glioma cells. AKT3 isoform may also be the key molecule in up-regulation and down-regulation of MMP13 via IL13. Required for the coordination of mitochondrial biogenesis with growth factor-induced increases in cellular energy demands. Down- regulation by RNA interference reduces the expression of the phosphorylated form of BAD,

resulting in the induction of caspase- dependent apoptosis.

Cellular Location

Nucleus. Cytoplasm. Membrane; Peripheral membrane protein
Note=Membrane-associated after cell stimulation leading to its translocation

Tissue Location

In adult tissues, it is highly expressed in brain, lung and kidney, but weakly in heart, testis and liver. In fetal tissues, it is highly expressed in heart, liver and brain and not at all in kidney

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.