

KD-Validated Anti-Diacylglycerol O-acyltransferase 1 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody
Catalog # AGI1503

Product Information

Application	WB, FC
Primary Accession	O75907
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB5155
Calculated MW	55278
Gene Name	DGAT1
Aliases	DGAT1; Diacylglycerol O-Acyltransferase 1; ARGP1; DGAT; Acyl-CoA Retinol O-Fatty-Acyltransferase; Diglyceride Acyltransferase; EC 2.3.1.20; ARAT; Acyl Coenzyme A:Cholesterol Acyltransferase Related Gene 1; Diacylglycerol O-Acyltransferase (Mouse) Homolog; Acyl-CoA:Diacylglycerol Acyltransferase; Retinol O-Fatty-Acyltransferase; ACAT Related Gene Product 1; ACAT-Related Gene Product 1; EC 2.3.1.76; EC 2.3.1; DIAR7; AGRP1
Immunogen	A synthesized peptide derived from human DGAT1

Additional Information

Gene ID	8694
Other Names	Diacylglycerol O-acyltransferase 1, 2.3.1.20, ACAT-related gene product 1, Acyl-CoA retinol O-fatty-acyltransferase, ARAT, Retinol O-fatty-acyltransferase, 2.3.1.76, Diglyceride acyltransferase, DGAT1 {ECO:0000303 PubMed:16214399, ECO:0000312 HGNC:HGNC:2843}

Protein Information

Name	DGAT1 {ECO:0000303 PubMed:16214399, ECO:0000312 HGNC:HGNC:2843}
Function	Catalyzes the terminal and only committed step in triacylglycerol synthesis by using diacylglycerol and fatty acyl CoA as substrates (PubMed: 16214399 , PubMed: 18768481 , PubMed: 28420705 , PubMed: 32433610 , PubMed: 32433611 , PubMed: 9756920). Highly expressed in epithelial cells of the small intestine and its activity is essential for the absorption of dietary fats (PubMed: 18768481). In liver, plays a role in esterifying exogenous fatty acids to glycerol, and is required to synthesize fat for storage (PubMed: 16214399). Also present in female mammary glands, where it produces fat in the milk (By similarity). May be involved in VLDL (very low density lipoprotein) assembly (PubMed: 18768481). In contrast to DGAT2 it is not essential for survival (By similarity). Functions as the major acyl-CoA retinol acyltransferase (ARAT) in

the skin, where it acts to maintain retinoid homeostasis and prevent retinoid toxicity leading to skin and hair disorders (PubMed:[16214399](#)). Exhibits additional acyltransferase activities, including acyl CoA:monoacylglycerol acyltransferase (MGAT), wax monoester and wax diester synthases (By similarity). Also able to use 1-monoalkylglycerol (1-MAkG) as an acyl acceptor for the synthesis of monoalkyl-monoacylglycerol (MAMAG) (PubMed:[28420705](#)).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q9Z2A7};
Multi-pass membrane protein

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.