10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

KD-Validated Anti-MYD88 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody
Catalog # AGI1591

Product Information

Application	WB, FC, ICC
Primary Accession	Q99836
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	24GB240
Calculated MW	33233
Gene Name	MYD88
Aliases	MYD88 Innate Immune Signal Transduction Adaptor; Myeloid Differentiation Primary Response Protein MyD88; Myeloid Differentiation Primary Response Gene (88); Myeloid Differentiation Primary Response 88; TLR Adaptor MYD88; Mutant Myeloid Differentiation Primary Response 88; MYD88D; IMD68; WM1
Immunogen	A synthesized peptide derived from human MyD88

Additional Information

Gene ID	4615
Other Names	Myeloid differentiation primary response protein MyD88, MYD88 (HGNC:7562)

Protein Information

Name	MYD88 (HGNC:7562)
Function	Adapter protein involved in the Toll-like receptor and IL-1 receptor signaling pathway in the innate immune response (PubMed:15361868, PubMed:18292575, PubMed:33718825, PubMed:37971847). Acts via IRAK1, IRAK2, IRF7 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:15361868, PubMed:19506249, PubMed:24316379, PubMed:40638072). Increases IL-8 transcription (PubMed:9013863). Involved in IL-18- mediated signaling pathway. Activates IRF1 resulting in its rapid migration into the nucleus to mediate an efficient induction of IFN- beta, NOS2/INOS, and IL12A genes. Upon TLR8 activation by GU-rich single-stranded RNA (GU-rich RNA) derived from viruses such as SARS- CoV-2, SARS-CoV and HIV-1, induces IL1B release through NLRP3 inflammasome activation (PubMed:33718825). MyD88-mediated signaling in intestinal epithelial cells is crucial for maintenance of gut homeostasis and controls the expression of the antimicrobial lectin REG3G in the small intestine (By similarity).
Cellular Location	Cytoplasm. Nucleus

Tissue Location	Ubiquitous..

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.