

KD-Validated Anti-DGCR8 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1595

Product Information

Application	WB, ICC
Primary Accession	Q8WYQ5
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB1615
Calculated MW	86045
Gene Name	DGCR8
Aliases	DGCR8; DGCR8 Microprocessor Complex Subunit; DGCRK6; DiGeorge Syndrome Critical Region Gene 8; Microprocessor Complex Subunit DGCR8; DiGeorge Syndrome Critical Region 8; C22orf12; Pasha; Gy1; DGCR8, Microprocessor Complex Subunit; Chromosome 22 Open Reading Frame 12; C22ORF12; PASHA; GY1
Immunogen	A synthesized peptide derived from human DGCR8

Additional Information

Gene ID	54487
Other Names	Microprocessor complex subunit DGCR8, DiGeorge syndrome critical region 8, DGCR8, C22orf12, DGCRK6

Protein Information

Name	DGCR8
Synonyms	C22orf12, DGCRK6
Function	Component of the microprocessor complex that acts as a RNA- and heme-binding protein that is involved in the initial step of microRNA (miRNA) biogenesis. Component of the microprocessor complex that is required to process primary miRNA transcripts (pri-miRNAs) to release precursor miRNA (pre-miRNA) in the nucleus. Within the microprocessor complex, DGCR8 function as a molecular anchor necessary for the recognition of pri-miRNA at dsRNA-ssRNA junction and directs DROSHA to cleave 11 bp away from the junction to release hairpin-shaped pre-miRNAs that are subsequently cut by the cytoplasmic DICER to generate mature miRNAs (PubMed: 26027739 , PubMed: 26748718). The heme- bound DGCR8 dimer binds pri-miRNAs as a cooperative trimer (of dimers) and is active in triggering pri-miRNA cleavage, whereas the heme-free DGCR8 monomer binds pri-miRNAs as a dimer and is much less active. Both double-stranded and single-stranded regions of a pri-miRNA are required for its binding (PubMed: 15531877 ,

PubMed:[15574589](#), PubMed:[15589161](#), PubMed:[16751099](#), PubMed:[16906129](#), PubMed:[16963499](#), PubMed:[17159994](#)). Specifically recognizes and binds N6-methyladenosine (m6A)-containing pri-miRNAs, a modification required for pri-miRNAs processing (PubMed:[25799998](#)). Involved in the silencing of embryonic stem cell self-renewal (By similarity). Also plays a role in DNA repair by promoting the recruitment of RNF168 to RNF8 and MDC1 at DNA double- strand breaks and subsequently the clearance of DNA breaks (PubMed:[34188037](#)).

Cellular Location

Nucleus. Nucleus, nucleolus. Note=Colocalizes with nucleolin and DROSHA in the nucleolus. Mostly detected in the nucleolus as electron-dense granular patches around the fibrillar center (FC) and granular component (GC). Also detected in the nucleoplasm as small foci adjacent to splicing speckles near the chromatin structure. Localized with DROSHA in GW bodies (GWBs), also known as P-bodies (PubMed:17159994)

Tissue Location

Ubiquitously expressed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.