

KD-Validated Anti-Peptidylprolyl Isomerase A Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1635

Product Information

Application	WB, FC, ICC
Primary Accession	P62937
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	24GB905
Calculated MW	18012
Gene Name	PPIA
Aliases	Peptidylprolyl Isomerase A; CYPA; Peptidyl-Prolyl Cis-Trans Isomerase A; Cyclosporin A-Binding Protein; Cyclophilin A; Rotamase A; EC 5.2.1.8; PPIase A; Epididymis Secretory Sperm Binding Protein Li 69p; Peptidylprolyl Isomerase A (Cyclophilin A); T Cell Cyclophilin; HEL-S-69p; CYPH
Immunogen	A synthesized peptide derived from human Cyclophilin A

Additional Information

Gene ID	5478
Other Names	Peptidyl-prolyl cis-trans isomerase A, PPIase A, 5.2.1.8, Cyclophilin A, Cyclosporin A-binding protein, Rotamase A, Peptidyl-prolyl cis-trans isomerase A, N-terminally processed, PPIA, CYPA

Protein Information

Name	PPIA
Synonyms	CYPA
Function	Catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides (PubMed: 2001362 , PubMed: 20676357 , PubMed: 21245143 , PubMed: 21593166 , PubMed: 25678563). Exerts a strong chemotactic effect on leukocytes partly through activation of one of its membrane receptors BSG/CD147, initiating a signaling cascade that culminates in MAPK/ERK activation (PubMed: 11943775 , PubMed: 21245143). Activates endothelial cells (ECs) in a pro-inflammatory manner by stimulating activation of NF-kappa-B and ERK, JNK and p38 MAP-kinases and by inducing expression of adhesion molecules including SELE and VCAM1 (PubMed: 15130913). Induces apoptosis in ECs by promoting the FOXO1-dependent expression of CCL2 and BCL2L11 which are involved in EC chemotaxis and apoptosis (PubMed: 31063815). In response to oxidative stress, initiates proapoptotic and antiapoptotic signaling

in ECs via activation of NF-kappa-B and AKT1 and up-regulation of antiapoptotic protein BCL2 (PubMed:[23180369](#)). Negatively regulates MAP3K5/ASK1 kinase activity, autophosphorylation and oxidative stress-induced apoptosis mediated by MAP3K5/ASK1 (PubMed:[26095851](#)). Necessary for the assembly of TARDBP in heterogeneous nuclear ribonucleoprotein (hnRNP) complexes and regulates TARDBP binding to RNA UG repeats and TARDBP-dependent expression of HDAC6, ATG7 and VCP which are involved in clearance of protein aggregates (PubMed:[25678563](#)). Plays an important role in platelet activation and aggregation (By similarity). Regulates calcium mobilization and integrin ITGA2B:ITGB3 bidirectional signaling via increased ROS production as well as by facilitating the interaction between integrin and the cell cytoskeleton (By similarity). Binds heparan sulfate glycosaminoglycans (PubMed:[11943775](#)). Inhibits replication of influenza A virus (IAV) (PubMed:[19207730](#)). Inhibits ITCH/AIP4-mediated ubiquitination of matrix protein 1 (M1) of IAV by impairing the interaction of ITCH/AIP4 with M1, followed by the suppression of the nuclear export of M1, and finally reduction of the replication of IAV (PubMed:[22347431](#), PubMed:[30328013](#)).

Cellular Location

Cytoplasm. Secreted Nucleus Cell membrane. Note=Secretion occurs in response to oxidative stress in vascular smooth muscle through a vesicular secretory pathway that includes Rho GTPase signaling, actin remodeling, and myosin II activation.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.