

KD-Validated Anti-TNF Alpha Induced Protein 8 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1726

Product Information

Application	WB, FC
Primary Accession	O95379
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	24GB2315
Calculated MW	23003
Gene Name	TNFAIP8
Aliases	TNFAIP8; TNF Alpha Induced Protein 8; MDC-3.13; SCC-S2; GG2-1; Head And Neck Tumor And Metastasis-Related Protein; Tumor Necrosis Factor Alpha-Induced Protein 8; NF-Kappa-B-Inducible DED-Containing Protein; TNF-Induced Protein GG2-1; NDED; Tumor Necrosis Factor, Alpha-Induced Protein 8; Tumor Necrosis Factor, Alpha Induced Protein 8; TNF Alpha-Induced Protein 8; SCCS2
Immunogen	A synthesized peptide derived from human TNFAIP8

Additional Information

Gene ID	25816
Other Names	Tumor necrosis factor alpha-induced protein 8, TNF alpha-induced protein 8, Head and neck tumor and metastasis-related protein, MDC-3.13, NF-kappa-B-inducible DED-containing protein, NDED, SCC-S2, TNF-induced protein GG2-1, TNFAIP8

Protein Information

Name	TNFAIP8
Function	Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.
Cellular Location	Cytoplasm.
Tissue Location	Expressed at high levels in the spleen, lymph node, thymus, thyroid, bone marrow and placenta. Expressed at high levels both in various tumor tissues, unstimulated and cytokine-activated cultured cells. Expressed at low levels in the spinal cord, ovary, lung, adrenal glands, heart, brain, testis and skeletal

muscle

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.