

KD-Validated Anti-UBR5 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1756

Product Information

Application	WB, FC, ICC
Primary Accession	O95071
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	24GB2540
Calculated MW	309352
Gene Name	UBR5
Aliases	UBR5; Ubiquitin Protein Ligase E3 Component N-Recognin 5; HYD; EDD; KIAA0896; EDD1; DD5; E3 Ubiquitin-Protein Ligase, HECT Domain-Containing 1; HECT-Type E3 Ubiquitin Transferase UBR5; Hyperplastic Discs Protein Homolog; E3 Ubiquitin-Protein Ligase UBR5; Progesterone-Induced Protein; E3 Ubiquitin Protein Ligase, HECT Domain Containing, 1; E3 Identified By Differential Display; EC 2.3.2.26; EC 6.3.2; HHYD
Immunogen	A synthesized peptide derived from human EDD

Additional Information

Gene ID	51366
Other Names	E3 ubiquitin-protein ligase UBR5, 2.3.2.26, E3 ubiquitin-protein ligase, HECT domain-containing 1, Hyperplastic discs protein homolog, hHYD, Progesterone-induced protein, UBR5

Protein Information

Name	UBR5
Function	E3 ubiquitin-protein ligase involved in different protein quality control pathways in the cytoplasm and nucleus (PubMed: 29033132 , PubMed: 33208877 , PubMed: 37478846 , PubMed: 37478862). Mainly acts as a ubiquitin chain elongator that extends pre-ubiquitinated substrates (PubMed: 29033132 , PubMed: 37409633). Component of the N-end rule pathway: ubiquitinates proteins bearing specific N-terminal residues that are destabilizing according to the N-end rule, leading to their degradation (By similarity). Recognizes type-1 N-degrons, containing positively charged amino acids (Arg, Lys and His) (By similarity). Together with UBR4, part of a cytoplasm protein quality control pathway that prevents protein aggregation by catalyzing assembly of heterotypic 'Lys-11'-/'Lys-48'-linked branched ubiquitin chains on aggregated proteins, leading to substrate recognition by the segregase p97/VCP and degradation by the proteasome: UBR5 is probably branching multiple 'Lys-48'-linked chains of substrates initially modified with

mixed conjugates by UBR4 (PubMed:[29033132](#)). Together with ITCH, catalyzes 'Lys-48'-'Lys-63'-branched ubiquitination of TXNIP, leading to its degradation: UBR5 mediates branching of 'Lys-48'-linked chains of substrates initially modified with 'Lys-63'-linked conjugates by ITCH (PubMed:[29378950](#)). Catalytic component of a nuclear protein quality control pathway that mediates ubiquitination and degradation of unpaired transcription factors (i.e. transcription factors that are not assembled into functional multiprotein complexes): specifically recognizes and binds degrons that are not accessible when transcription regulators are associated with their coactivators (PubMed:[37478846](#), PubMed:[37478862](#)). Ubiquitinates various unpaired transcription regulator (MYC, SUPT4H1, SUPT5H, CDC20 and MCRS1), as well as ligand- bound nuclear receptors (ESR1, NR1H3, NR3C1, PGR, RARA, RXRA AND VDR) that are not associated with their nuclear receptor coactivators (NCOAs) (PubMed:[33208877](#), PubMed:[37478846](#), PubMed:[37478862](#)). Involved in maturation and/or transcriptional regulation of mRNA by mediating polyubiquitination and activation of CDK9 (PubMed:[21127351](#)). Also acts as a regulator of DNA damage response by acting as a suppressor of RNF168, an E3 ubiquitin-protein ligase that promotes accumulation of 'Lys-63'-linked histone H2A and H2AX at DNA damage sites, thereby acting as a guard against excessive spreading of ubiquitinated chromatin at damaged chromosomes (PubMed:[22884692](#)). Regulates DNA topoisomerase II binding protein (TopBP1) in the DNA damage response (PubMed:[11714696](#)). Ubiquitinates acetylated PCK1 (PubMed:[21726808](#)). Acts as a positive regulator of the canonical Wnt signaling pathway by mediating (1) ubiquitination and stabilization of CTNNB1, and (2) 'Lys- 48'-linked ubiquitination and degradation of TLE3 (PubMed:[21118991](#), PubMed:[28689657](#)). Promotes disassembly of the mitotic checkpoint complex (MCC) from the APC/C complex by catalyzing ubiquitination of BUB1B, BUB3 and CDC20 (PubMed:[35217622](#)). Plays an essential role in extraembryonic development (By similarity). Required for the maintenance of skeletal tissue homeostasis by acting as an inhibitor of hedgehog (HH) signaling (By similarity).

Cellular Location

Nucleus. Cytoplasm

Tissue Location

Widely expressed. Most abundant in testis and expressed at high levels in brain, pituitary and kidney

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.