

KD-Validated Anti-Mitotic Arrest Deficient 2 Like 2 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1870

Product Information

Application	WB, FC
Primary Accession	Q9UI95
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	24GB4950
Calculated MW	24334
Gene Name	MAD2L2
Aliases	MAD2L2; Mitotic Arrest Deficient 2 Like 2; MAD2B; REV7; POLZ2; FANCV; MAD2 (Mitotic Arrest Deficient, Yeast, Homolog)-Like 2; Polymerase (DNA-Directed), Zeta 2, Accessory Subunit; Mitotic Spindle Assembly Checkpoint Protein MAD2B; Mitotic Arrest Deficient 2-Like Protein 2; Mitotic Arrest Deficient Homolog-Like 2; MAD2-Like Protein 2; REV7 Homolog; hREV7; MAD2 Mitotic Arrest Deficient-Like 2
Immunogen	A synthesized peptide derived from human Mad2L2

Additional Information

Gene ID	10459
Other Names	Mitotic spindle assembly checkpoint protein MAD2B, Mitotic arrest deficient 2-like protein 2, MAD2-like protein 2, REV7 homolog, hREV7, MAD2L2, MAD2B, REV7

Protein Information

Name	MAD2L2
Synonyms	MAD2B, REV7
Function	Adapter protein able to interact with different proteins and involved in different biological processes (PubMed: 11459825 , PubMed: 11459826 , PubMed: 17296730 , PubMed: 17719540 , PubMed: 19443654 , PubMed: 29656893). Mediates the interaction between the error-prone DNA polymerase zeta catalytic subunit REV3L and the inserter polymerase REV1, thereby mediating the second polymerase switching in translesion DNA synthesis (PubMed: 20164194 , PubMed: 23143872). Translesion DNA synthesis releases the replication blockade of replicative polymerases, stalled in presence of DNA lesions (PubMed: 20164194). Component of the shieldin complex, which plays an important role in repair of DNA double-stranded

breaks (DSBs) (PubMed:[29656893](#)). During G1 and S phase of the cell cycle, the complex functions downstream of TP53BP1 to promote non-homologous end joining (NHEJ) and suppress DNA end resection (PubMed:[29656893](#)). Mediates various NHEJ-dependent processes including immunoglobulin class-switch recombination, and fusion of unprotected telomeres (PubMed:[29656893](#)). May also regulate another aspect of cellular response to DNA damage through regulation of the JNK-mediated phosphorylation and activation of the transcriptional activator ELK1 (PubMed:[17296730](#)). Inhibits the FZR1- and probably CDC20-mediated activation of the anaphase promoting complex APC thereby regulating progression through the cell cycle (PubMed:[11459825](#), PubMed:[17719540](#)). Regulates TCF7L2-mediated gene transcription and may play a role in epithelial-mesenchymal transdifferentiation (PubMed:[19443654](#)).

Cellular Location

Nucleus. Cytoplasm, cytoskeleton, spindle. Cytoplasm. Chromosome.
Note=Recruited to sites of chromosomal double-stranded breaks during G1 and S phase of the cell cycle

Tissue Location

Ubiquitously expressed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.