

KD-Validated Anti-Phospho-NDRG1 (Ser330) Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1873

Product Information

Application	WB, FC
Primary Accession	Q92597
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	24GB4985
Calculated MW	42835
Gene Name	NDRG1
Aliases	NDRG1; N-Myc Downstream Regulated 1; DRG1; RTP; CAP43; TDD5; NDR1; Reducing Agents And Tunicamycin-Responsive Protein; N-Myc Downstream-Regulated Gene 1 Protein; Nickel-Specific Induction Protein Cap43; Differentiation-Related Gene 1 Protein; Protein NDRG1; DRG-1; Protein Regulated By Oxygen-1; PROXY1; CMT4D; HMSNL; RIT42; TARG1; Rit42; NMSL; GC4
Immunogen	A synthesized peptide derived from human Phospho-NDRG1 (Ser330)

Additional Information

Gene ID	10397
Other Names	Protein NDRG1, Differentiation-related gene 1 protein, DRG-1, N-myc downstream-regulated gene 1 protein, Nickel-specific induction protein Cap43, Reducing agents and tunicamycin-responsive protein, RTP, Rit42, NDRG1, CAP43, DRG1, RTP

Protein Information

Name	NDRG1
Synonyms	CAP43, DRG1, RTP
Function	Stress-responsive protein involved in hormone responses, cell growth, and differentiation. Acts as a tumor suppressor in many cell types. Necessary but not sufficient for p53/TP53-mediated caspase activation and apoptosis. Has a role in cell trafficking, notably of the Schwann cell, and is necessary for the maintenance and development of the peripheral nerve myelin sheath. Required for vesicular recycling of CDH1 and TF. May also function in lipid trafficking. Protects cells from spindle disruption damage. Functions in p53/TP53-dependent mitotic spindle checkpoint. Regulates microtubule dynamics and maintains euploidy.

Cellular Location

Cytoplasm, cytosol. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cell membrane Note=Mainly cytoplasmic but differentially localized to other regions Associates with the plasma membrane in intestinal epithelia and lactating mammary gland. Translocated to the nucleus in a p53/TP53- dependent manner. In prostate epithelium and placental chorion, located in both the cytoplasm and in the nucleus. No nuclear localization in colon epithelium cells. In intestinal mucosa, prostate and renal cortex, located predominantly adjacent to adherens junctions Cytoplasmic with granular staining in proximal tubular cells of the kidney and salivary gland ducts. Recruits to the membrane of recycling/sorting and late endosomes via binding to phosphatidylinositol 4-phosphate. Associates with microtubules Colocalizes with TUBG1 in the centrosome. Cytoplasmic location increased with hypoxia. Phosphorylated form found associated with centromeres during S-phase of mitosis and with the plasma membrane

Tissue Location

Ubiquitous; expressed most prominently in placental membranes and prostate, kidney, small intestine, and ovary tissues Also expressed in heart, brain, skeletal muscle, lung, liver and pancreas. Low levels in peripheral blood leukocytes and in tissues of the immune system. Expressed mainly in epithelial cells. Also found in Schwann cells of peripheral neurons. Reduced expression in adenocarcinomas compared to normal tissues. In colon, prostate and placental membranes, the cells that border the lumen show the highest expression.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.