

KD-Validated Anti-Catenin Delta 1 Mouse Monoclonal Antibody

Mouse monoclonal antibody
Catalog # AGI1917

Product Information

Application	WB
Primary Accession	O60716
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1
Clone Names	24GB5775
Calculated MW	108170
Gene Name	CTNND1
Aliases	CTNND1; Catenin Delta 1; KIAA0384; P120; Catenin (Cadherin-Associated Protein), Delta 1; Cadherin-Associated Src Substrate; Catenin Delta-1; P120 Catenin; P120CAS; P120CTN; CTNND; CAS; P120(CAS); P120(CTN); P120(Ctn); P120(Cas); P120cas; P120ctn; BCDS2
Immunogen	Recombinant protein of human CTNND1

Additional Information

Gene ID	1500
Other Names	Catenin delta-1, Cadherin-associated Src substrate, CAS, p120 catenin, p120(ctn), p120(cas), CTNND1 (HGNC:2515), KIAA0384

Protein Information

Name	CTNND1 (HGNC:2515)
Synonyms	KIAA0384
Function	Key regulator of cell-cell adhesion that associates with and regulates the cell adhesion properties of both C-, E- and N-cadherins, being critical for their surface stability (PubMed: 14610055 , PubMed: 20371349). Promotes localization and retention of DSG3 at cell- cell junctions, via its interaction with DSG3 (PubMed: 18343367). Beside cell-cell adhesion, regulates gene transcription through several transcription factors including ZBTB33/Kaiso2 and GLIS2, and the activity of Rho family GTPases and downstream cytoskeletal dynamics (PubMed: 10207085 , PubMed: 20371349). Implicated both in cell transformation by SRC and in ligand-induced receptor signaling through the EGF, PDGF, CSF-1 and ERBB2 receptors (PubMed: 17344476).
Cellular Location	Cell junction, adherens junction. Cytoplasm. Nucleus. Cell membrane. Cell junction. Note=Interaction with GLIS2 promotes nuclear translocation (By

similarity). Detected at cell-cell contacts (PubMed:15240885, PubMed:17047063). NANOS1 induces its translocation from sites of cell-cell contact to the cytoplasm (PubMed:17047063). CDH1 enhances cell membrane localization (PubMed:15240885). Localizes to cell-cell contacts as keratinocyte differentiation progresses (By similarity) {ECO:0000250 | UniProtKB:P30999, ECO:0000269 | PubMed:11896187, ECO:0000269 | PubMed:15240885, ECO:0000269 | PubMed:17047063} [Isoform 2A]: Nucleus [Isoform 4A]: Cytoplasm

Tissue Location

Expressed in vascular endothelium. Melanocytes and melanoma cells primarily express the long isoform 1A, whereas keratinocytes express shorter isoforms, especially 3A. The shortest isoform 4A, is detected in normal keratinocytes and melanocytes, and generally lost from cells derived from squamous cell carcinomas or melanomas. The C-terminal alternatively spliced exon B is present in the p120ctn transcripts in the colon, intestine and prostate, but lost in several tumor tissues derived from these organs

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.