

KD-Validated Anti-GRK2 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI1966

Product Information

Application	WB, FC, ICC
Primary Accession	P25098
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	23GB1515
Calculated MW	79574
Gene Name	GRK2
Aliases	GRK2; G Protein-Coupled Receptor Kinase 2; BARK1; ADRBK1; Beta-Adrenergic Receptor Kinase 1; EC 2.7.11.15; Beta-ARK-1; Adrenergic, Beta, Receptor Kinase 1; G-Protein Coupled Receptor Kinase; Adrenergic Beta Receptor Kinase 1; BETA-ARK1; EC 2.7.11; BARK
Immunogen	A synthesized peptide derived from human GRK2

Additional Information

Gene ID	156
Other Names	Beta-adrenergic receptor kinase 1, Beta-ARK-1, 2.7.11.15, G-protein coupled receptor kinase 2 {ECO:0000312 HGNC:HGNC:289}, GRK2 (HGNC:289), ADRBK1, BARK, BARK1

Protein Information

Name	GRK2 {ECO:0000303 PubMed:40384633, ECO:0000312 HGNC:HGNC:289}
Function	Specifically phosphorylates the activated forms of G protein- coupled receptors (PubMed: 14976224 , PubMed: 16179383 , PubMed: 19306925 , PubMed: 19715378 , PubMed: 21799898 , PubMed: 23077507 , PubMed: 30284927 , PubMed: 39138140 , PubMed: 40384633). Such receptor phosphorylation initiates beta-arrestin-mediated receptor internalization, and signaling events leading to their desensitization (PubMed: 14976224 , PubMed: 16179383 , PubMed: 19306925 , PubMed: 19715378 , PubMed: 21799898 , PubMed: 23077507 , PubMed: 30284927 , PubMed: 39138140 , PubMed: 40384633). Phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors, probably inducing a desensitization of them (PubMed: 19715378). Phosphorylates catecholamine-activated ADRB2 to regulate physiological cardiomyocyte contraction rate responses (By similarity). Also phosphorylates ligand-bound C3a and C5a anaphylatoxin receptors (C3AR1 and C5AR1, respectively), leading to receptor desensitization (PubMed: 21799898 , PubMed: 23077507). Positively regulates ciliary smoothened (SMO) signaling pathway by mediating

phosphorylation of SMO, facilitating SMO trafficking into the cilium and the stimulation of SMO activity (PubMed:[39138140](#)). Also positively regulates SMO signaling by mediating phosphorylation of GPR161, a key inhibitor SMO signaling, promoting ciliary export of GPR161 (PubMed:[40384633](#)). Key regulator of LPAR1 signaling (PubMed:[19306925](#)). Competes with RALA for binding to LPAR1 thus affecting the signaling properties of the receptor (PubMed:[19306925](#)). Desensitizes LPAR1 and LPAR2 in a phosphorylation-independent manner (PubMed:[19306925](#)). Inhibits relaxation of airway smooth muscle in response to blue light (PubMed:[30284927](#)).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P26817}. Cell membrane {ECO:0000250|UniProtKB:P21146}. Cell projection, cilium membrane. Postsynapse {ECO:0000250|UniProtKB:P26817}. Presynapse {ECO:0000250|UniProtKB:P26817}

Tissue Location

Expressed in peripheral blood leukocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.