

KD-Validated Anti-Sp1 Transcription Factor Mouse Monoclonal Antibody

Mouse monoclonal antibody

Catalog # AGI1991

Product Information

Application	WB, FC, ICC
Primary Accession	P08047
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1 kappa
Clone Names	24GB7095
Calculated MW	80693
Gene Name	SP1
Aliases	SP1; Sp1 Transcription Factor; Transcription Factor Sp1; Specificity Protein 1; TSFP1
Immunogen	Recombinant protein of human SP1

Additional Information

Gene ID	6667
Other Names	Transcription factor Sp1, SP1, TSFP1

Protein Information

Name	SP1
Synonyms	TSFP1
Function	Transcription factor that can activate or repress transcription in response to physiological and pathological stimuli. Binds with high affinity to GC-rich motifs and regulates the expression of a large number of genes involved in a variety of processes such as cell growth, apoptosis, differentiation and immune responses. Highly regulated by post-translational modifications (phosphorylations, sumoylation, proteolytic cleavage, glycosylation and acetylation). Also binds the PDGFR-alpha G-box promoter. May have a role in modulating the cellular response to DNA damage. Implicated in chromatin remodeling. Plays an essential role in the regulation of FE65 gene expression. In complex with ATF7IP, maintains telomerase activity in cancer cells by inducing TERT and TERC gene expression. Isoform 3 is a stronger activator of transcription than isoform 1. Positively regulates the transcription of the core clock component BMAL1 (PubMed:10391891, PubMed:11371615, PubMed:11904305, PubMed:14593115, PubMed:16377629, PubMed:16478997, PubMed:16943418, PubMed:17049555, PubMed:18171990, PubMed:18199680, PubMed:18239466,

PubMed:[18513490](#), PubMed:[18619531](#), PubMed:[19193796](#), PubMed:[20091743](#), PubMed:[21046154](#), PubMed:[21798247](#)). Plays a role in the recruitment of SMARCA4/BRG1 on the c-FOS promoter. Plays a role in protecting cells against oxidative stress following brain injury by regulating the expression of RNF112 (By similarity).

Cellular Location

Nucleus. Cytoplasm. Note=Nuclear location is governed by glycosylated/phosphorylated states. Insulin promotes nuclear location, while glucagon favors cytoplasmic location

Tissue Location

Up-regulated in adenocarcinomas of the stomach (at protein level). Isoform 3 is ubiquitously expressed at low levels

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.