

KD-Validated Anti-XIAP Mouse Monoclonal Antibody

Mouse monoclonal antibody

Catalog # AGI2004

Product Information

Application	WB
Primary Accession	P98170
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1,k
Clone Names	24GB8570
Calculated MW	56685
Gene Name	XIAP
Aliases	XIAP; X-Linked Inhibitor Of Apoptosis; BIRC4; HILP; API3; X-Linked Inhibitor Of Apoptosis, E3 Ubiquitin Protein Ligase; Baculoviral IAP Repeat-Containing Protein 4; RING-Type E3 Ubiquitin Transferase XIAP; E3 Ubiquitin-Protein Ligase XIAP; Inhibitor Of Apoptosis Protein 3; IAP-Like Protein 1; X-Linked IAP; HIAP-3; ILP-1; IAP-3; HIAP3; X-Linked Inhibitor Of Apoptosis Protein; Baculoviral IAP Repeat-Containing 4; IAP-Like Protein; EC 2.3.2.27; ILP1; MIHA; XLP2; IAP3; ILP
Immunogen	Recombinant human XIAP

Additional Information

Gene ID	331
Other Names	E3 ubiquitin-protein ligase XIAP, 2.3.2.27, Baculoviral IAP repeat-containing protein 4, IAP-like protein, ILP, hILP, Inhibitor of apoptosis protein 3, IAP-3, hIAP-3, hIAP3, RING-type E3 ubiquitin transferase XIAP, X-linked inhibitor of apoptosis protein, X-linked IAP, XIAP {ECO:0000303 PubMed:12121969, ECO:0000312 HGNC:HGNC:592}

Protein Information

Name	XIAP {ECO:0000303 PubMed:12121969, ECO:0000312 HGNC:HGNC:592}
Function	Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, copper homeostasis, mitogenic kinase signaling, cell proliferation, as well as cell invasion and metastasis (PubMed: 11257230 , PubMed: 11257231 , PubMed: 11447297 , PubMed: 12121969 , PubMed: 12620238 , PubMed: 17560374 , PubMed: 17967870 , PubMed: 19473982 , PubMed: 20154138 , PubMed: 22103349 , PubMed: 9230442). Acts as a direct caspase inhibitor (PubMed: 11257230 , PubMed: 11257231 , PubMed: 12620238). Directly bind to the active site pocket of CASP3 and CASP7 and obstructs substrate entry (PubMed: 11257230 , PubMed: 11257231 , PubMed: 16352606 , PubMed: 16916640). Inactivates CASP9 by keeping it in a monomeric, inactive

state (PubMed:[12620238](#)). Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and the target proteins for its E3 ubiquitin- protein ligase activity include: RIPK1, RIPK2, MAP3K2/MEKK2, DIABLO/SMAC, AIFM1, CCS, PTEN and BIRC5/survivin (PubMed:[17560374](#), PubMed:[17967870](#), PubMed:[19473982](#), PubMed:[20154138](#), PubMed:[22103349](#), PubMed:[22607974](#), PubMed:[29452636](#), PubMed:[30026309](#)). Acts as an important regulator of innate immunity by mediating 'Lys-63'-linked polyubiquitination of RIPK2 downstream of NOD1 and NOD2, thereby transforming RIPK2 into a scaffolding protein for downstream effectors, ultimately leading to activation of the NF-kappa-B and MAP kinases signaling (PubMed:[19667203](#), PubMed:[22607974](#), PubMed:[29452636](#), PubMed:[30026309](#)). 'Lys-63'-linked polyubiquitination of RIPK2 also promotes recruitment of the LUBAC complex to RIPK2 (PubMed:[22607974](#), PubMed:[29452636](#)). Regulates the BMP signaling pathway and the SMAD and MAP3K7/TAK1 dependent pathways leading to NF-kappa-B and JNK activation (PubMed:[17560374](#)). Ubiquitination of CCS leads to enhancement of its chaperone activity toward its physiologic target, SOD1, rather than proteasomal degradation (PubMed:[20154138](#)). Ubiquitination of MAP3K2/MEKK2 and AIFM1 does not lead to proteasomal degradation (PubMed:[17967870](#), PubMed:[22103349](#)). Plays a role in copper homeostasis by ubiquitinating COMMD1 and promoting its proteasomal degradation (PubMed:[14685266](#)). Can also function as E3 ubiquitin-protein ligase of the NEDD8 conjugation pathway, targeting effector caspases for neddylation and inactivation (PubMed:[21145488](#)). Ubiquitinates and therefore mediates the proteasomal degradation of BCL2 in response to apoptosis (PubMed:[29020630](#)). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the capability to kill cancer cells in a caspase-dependent and caspase- independent manner (PubMed:[22095281](#)). Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8 (PubMed:[22095281](#)). Acts as a positive regulator of Wnt signaling and ubiquitinates TLE1, TLE2, TLE3, TLE4 and AES (PubMed:[22304967](#)). Ubiquitination of TLE3 results in inhibition of its interaction with TCF7L2/TCF4 thereby allowing efficient recruitment and binding of the transcriptional coactivator beta-catenin to TCF7L2/TCF4 that is required to initiate a Wnt-specific transcriptional program (PubMed:[22304967](#)).

Cellular Location

Cytoplasm. Nucleus. Note=TLE3 promotes its nuclear localization.

Tissue Location

Expressed in colonic crypts (at protein level) (PubMed:30389919). Ubiquitous, except peripheral blood leukocytes (PubMed:8654366).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.