10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

KD-Validated Anti-TP53BP1 Mouse Monoclonal Antibody

Mouse monoclonal antibody
Catalog # AGI2089

Product Information

Application	WB
Primary Accession	Q12888
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Mouse IgG2b
Clone Names	24GB13320
Calculated MW	213574
Gene Name	TP53BP1
Aliases	TP53BP1; Tumor Protein P53 Binding Protein 1; 53BP1; TDRD30; P202; TP53-Binding Protein 1; P53-Binding Protein 1; P53BP1; Tumor Suppressor P53-Binding Protein 1; Tumor Protein P53-Binding Protein, 1; Tumor Protein 53-Binding Protein, 1
Immunogen	Recombinant protein of human TP53BP1

Additional Information

Gene ID	7158
Other Names	TP53-binding protein 1, 53BP1, p53-binding protein 1, p53BP1, TP53BP1 (HGNC:11999)

Protein Information

Name	TP53BP1 (HGNC:11999)
Function	Double-strand break (DSB) repair protein involved in response to DNA damage, telomere dynamics and class-switch recombination (CSR) during antibody genesis (PubMed:12364621, PubMed:17190600, PubMed:21144835, PubMed:22553214, PubMed:23333306, PubMed:27153538, PubMed:28241136, PubMed:31135337, PubMed:37696958). Plays a key role in the repair of double-strand DNA breaks (DSBs) in response to DNA damage by promoting non-homologous end joining (NHEJ)-mediated repair of DSBs and specifically counteracting the function of the homologous recombination (HR) repair protein BRCA1 (PubMed:22553214, PubMed:23333306, PubMed:23727112, PubMed:27153538, PubMed:31135337). In response to DSBs, phosphorylation by ATM promotes interaction with RIF1 and dissociation from NUDT16L1/TIRR, leading to recruitment to DSBs sites (PubMed:28241136). Recruited to DSBs sites by recognizing and binding histone H2A monoubiquitinated at 'Lys-15' (H2AK15Ub) and histone H4 dimethylated at 'Lys-20' (H4K20me2), two histone marks that are present at DSBs sites (PubMed:17190600, PubMed:23760478, PubMed:27153538, PubMed:28241136). Required for immunoglobulin class- switch

recombination (CSR) during antibody genesis, a process that involves the generation of DNA DSBs (PubMed:23345425). Participates in the repair and the orientation of the broken DNA ends during CSR (By similarity). In contrast, it is not required for classic NHEJ and V(D)J recombination (By similarity). Promotes NHEJ of dysfunctional telomeres via interaction with PAXIP1 (PubMed:23727112).

Cellular Location

Nucleus. Chromosome. Chromosome, centromere, kinetochore {ECO:0000250|UniProtKB:P70399}. Note=Localizes to the nucleus in absence of DNA damage (PubMed:28241136). Following DNA damage, recruited to sites of DNA damage, such as double strand breaks (DSBs): recognizes and binds histone H2A monoubiquitinated at 'Lys-15' (H2AK15Ub) and histone H4 dimethylated at 'Lys-20' (H4K20me2), two histone marks that are present at DSBs sites (PubMed:17190600, PubMed:23333306, PubMed:23760478, PubMed:24703952, PubMed:28241136, PubMed:31135337, PubMed:37696958). Associated with kinetochores during mitosis (By similarity). {ECO:0000250|UniProtKB:P70399, ECO:0000269|PubMed:17190600, ECO:0000269|PubMed:23333306, ECO:0000269|PubMed:23760478, ECO:0000269|PubMed:28241136, ECO:0000269|PubMed:37696958}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.