

KD-Validated Anti-HDAC1 Rabbit Polyclonal Antibody

Rabbit polyclonal antibody

Catalog # AGI2123

Product Information

Application	WB
Primary Accession	Q13547
Reactivity	Rat, Human, Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	55103
Gene Name	HDAC1
Aliases	HDAC1; Histone Deacetylase 1; HD1; GON-10; RPD3L1; KDAC1; Protein Decrotonylase HDAC1; Protein Deacetylase HDAC1; EC 3.5.1.98; Reduced Potassium Dependency, Yeast Homolog-Like 1; EC 3.5.1.-; RPD3
Immunogen	A synthesized peptide derived from human Histone Deacetylase 1

Additional Information

Gene ID	3065
Other Names	Histone deacetylase 1, HD1, 3.5.1.98, Protein deacetylase HDAC1, 3.5.1.-, Protein deacylase HDAC1, 3.5.1.-, HDAC1 {ECO:0000303 PubMed:10846170, ECO:0000312 HGNC:HGNC:4852}

Protein Information

Name	HDAC1 {ECO:0000303 PubMed:10846170, ECO:0000312 HGNC:HGNC:4852}
Function	Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed:16762839, PubMed:17704056, PubMed:28497810). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed:16762839, PubMed:17704056). Histone deacetylases act via the formation of large multiprotein complexes (PubMed:16762839, PubMed:17704056). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed:16428440, PubMed:28977666). As part of the SIN3B complex is recruited downstream of the constitutively active genes transcriptional start sites through interaction with histones and mitigates histone acetylation and RNA polymerase II progression within transcribed regions contributing to the regulation of transcription (PubMed:21041482). Also functions as a deacetylase for non-histone targets, such as NR1D2, RELA, SP1, SP3, STAT3, ZNF76 and TSHZ3 (PubMed:12837748, PubMed:16285960, PubMed:16337145, PubMed:16478997, PubMed:17996965, PubMed:19343227). Deacetylates SP proteins, SP1 and SP3, and regulates their function (PubMed:12837748,

PubMed:[16478997](#)). Component of the BRG1-RB1-HDAC1 complex, which negatively regulates the CREST-mediated transcription in resting neurons (PubMed:[19081374](#)). Upon calcium stimulation, HDAC1 is released from the complex and CREBBP is recruited, which facilitates transcriptional activation (PubMed:[19081374](#)). Deacetylates TSHZ3 and regulates its transcriptional repressor activity (PubMed:[19343227](#)). Deacetylates 'Lys-310' in RELA and thereby inhibits the transcriptional activity of NF-kappa-B (PubMed:[17000776](#)). Deacetylates NR1D2 and abrogates the effect of KAT5-mediated relieving of NR1D2 transcription repression activity (PubMed:[17996965](#)). Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development (By similarity). Involved in CIART-mediated transcriptional repression of the circadian transcriptional activator: CLOCK-BMAL1 heterodimer (By similarity). Required for the transcriptional repression of circadian target genes, such as PER1, mediated by the large PER complex or CRY1 through histone deacetylation (By similarity). In addition to protein deacetylase activity, also has protein-lysine deacylase activity: acts as a protein decrotonylase and delactylase by mediating decrotonylation ((2E)-butenoyl) and delactylation (lactoyl) of histones, respectively (PubMed:[28497810](#), PubMed:[35044827](#)).

Cellular Location

Nucleus

Tissue Location

Ubiquitous, with higher levels in heart, pancreas and testis, and lower levels in kidney and brain

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.