

KD-Validated Anti-KeRatin 17 Rabbit Polyclonal Antibody

Rabbit polyclonal antibody

Catalog # AGI2125

Product Information

Application	WB
Primary Accession	Q04695
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	48106
Gene Name	KRT17
Aliases	KRT17; KeRatin 17; KeRatin, Type I Cytoskeletal 17; KeRatin 17, Type I; CytokeRatin-17; CK-17; PCHC1; 39.1; K17; KeRatin-17; PC2; PC
Immunogen	A synthesized peptide derived from human Cytokeratin 17

Additional Information

Gene ID	3872
Other Names	Keratin, type I cytoskeletal 17, 39.1, Cytokeratin-17, CK-17, Keratin-17, K17, KRT17

Protein Information

Name	KRT17
Function	Type I keratin involved in the formation and maintenance of various skin appendages, specifically in determining shape and orientation of hair (By similarity). Required for the correct growth of hair follicles, in particular for the persistence of the anagen (growth) state (By similarity). Modulates the function of TNF in the specific context of hair cycling. Regulates protein synthesis and epithelial cell growth through binding to the adapter protein SFN and by stimulating Akt/mTOR pathway (By similarity). Involved in tissue repair. May be a marker of basal cell differentiation in complex epithelia and therefore indicative of a certain type of epithelial 'stem cells'. Acts as a promoter of epithelial proliferation by acting a regulator of immune response in skin: promotes Th1/Th17-dominated immune environment contributing to the development of basaloid skin tumors (By similarity). May act as an autoantigen in the immunopathogenesis of psoriasis, with certain peptide regions being a major target for autoreactive T-cells and hence causing their proliferation.
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q9QWL7}.
Tissue Location	Expressed in the outer root sheath and medulla region of hair follicle specifically from eyebrow and beard, digital pulp, nail matrix and nail bed

epithelium, mucosal stratified squamous epithelia and in basal cells of oral epithelium, palmoplantar epidermis and sweat and mammary glands. Also expressed in myoepithelium of prostate, basal layer of urinary bladder, cambial cells of sebaceous gland and in exocervix (at protein level)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.