

KD-Validated Anti-YWHAG Rabbit Polyclonal Antibody

Rabbit polyclonal antibody

Catalog # AGI2133

Product Information

Application	WB
Primary Accession	P61981
Reactivity	Rat, Human, Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	28303
Gene Name	YWHAG
Aliases	YWHAG; Tyrosine 3-Monooxygenase/Tryptophan 5-Monooxygenase Activation Protein Gamma; 14-3-3GAMMA; PPP1R170; Tyrosine 3-Monooxygenase/Tryptophan 5-Monooxygenase Activation Protein, Gamma Polypeptide; Protein Phosphatase 1, Regulatory Subunit 170; Protein Kinase C Inhibitor Protein 1; 14-3-3 Protein Gamma; KCIP-1; 14-3-3 Gamma; 14-3-3γ; 14-3-3G; EIEE56; DEE56
Immunogen	A synthesized peptide derived from human 14-3-3 gamma

Additional Information

Gene ID	7532
Other Names	14-3-3 protein gamma, Protein kinase C inhibitor protein 1, KCIP-1, 14-3-3 protein gamma, N-terminally processed, YWHAG (HGNC:12852)

Protein Information

Name	YWHAG (HGNC:12852)
Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed: 15696159 , PubMed: 16511572 , PubMed: 36732624). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed: 15696159 , PubMed: 16511572 , PubMed: 36732624). Binding generally results in the modulation of the activity of the binding partner (PubMed: 16511572). Promotes inactivation of WDR24 component of the GATOR2 complex by binding to phosphorylated WDR24 (PubMed: 36732624). Participates in the positive regulation of NMDA glutamate receptor activity by promoting the L- glutamate secretion through interaction with BEST1 (PubMed: 29121962). Reduces keratinocyte intercellular adhesion, via interacting with PKP1 and sequestering it in the cytoplasm, thereby reducing its incorporation into desmosomes (PubMed: 29678907). Plays a role in mitochondrial protein catabolic process (also named MALM) that promotes the degradation of damaged proteins inside mitochondria (PubMed: 22532927).

Cellular Location	Cytoplasm, cytosol. Mitochondrion matrix. Note=Translocates to the mitochondrial matrix following induction of MALM (mitochondrial protein catabolic process).
Tissue Location	Highly expressed in brain, skeletal muscle, and heart.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.