

KD-Validated Anti-KDM2A Rabbit Monoclonal Antibody

Rabbit monoclonal Antibody

Catalog # AGI2203

Product Information

Application	WB, FC
Primary Accession	Q9Y2K7
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	25GB160
Calculated MW	132793
Gene Name	KDM2A
Aliases	KDM2A; Lysine Demethylase 2A; JHDM1A; FBL11; CXXC8; FBL7; F-Box And Leucine-Rich Repeat Protein 11; KIAA1004; LILINA; FBXL11; JmjC Domain-Containing Histone Demethylation Protein 1A; Jumonji C Domain-Containing Histone Demethylase 1A; [Histone-H3]-Lysine-36 Demethylase 1A; Lysine (K)-Specific Demethylase 2A; CXXC-Type Zinc Finger Protein 8; Lysine-Specific Demethylase 2A; F-Box/LRR-Repeat Protein 11; DKFZP434M1735; EC 1.14.11.27; FLJ00115; F-Box Protein Lilina; F-Box Protein FBL11; F-Box Protein FBL7
Immunogen	A synthesized peptide derived from human FBXL11

Additional Information

Gene ID	22992
Other Names	Lysine-specific demethylase 2A, 1.14.11.27, CXXC-type zinc finger protein 8, F-box and leucine-rich repeat protein 11, F-box protein FBL7, F-box protein Lilina, F-box/LRR-repeat protein 11, JmjC domain-containing histone demethylation protein 1A, [Histone-H3]-lysine-36 demethylase 1A, KDM2A

Protein Information

Name	KDM2A
Function	Histone demethylase that specifically demethylates 'Lys-36' of histone H3, thereby playing a central role in histone code. Preferentially demethylates dimethylated H3 'Lys-36' residue while it has weak or no activity for mono- and tri-methylated H3 'Lys-36'. May also recognize and bind to some phosphorylated proteins and promote their ubiquitination and degradation. Required to maintain the heterochromatic state. Associates with centromeres and represses transcription of small non-coding RNAs that are encoded by the clusters of satellite repeats at the centromere. Required to sustain centromeric integrity and genomic stability, particularly during mitosis. Regulates circadian gene expression by repressing the transcriptional activator activity of CLOCK-BMAL1 heterodimer and RORA in a catalytically-

independent manner (PubMed:[26037310](#)).

Cellular Location

Nucleus, nucleoplasm. Chromosome Note=Punctate expression throughout the nucleoplasm and enriched in the perinucleolar region (PubMed:19001877, PubMed:20417597). Specifically nucleates at CpG islands where its presence results in chromatin depleted in H3K36me2 (PubMed:19001877, PubMed:20417597)

Tissue Location

Widely expressed, with highest levels in brain, testis and ovary, followed by lung.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.