

KD-Validated Anti-PKN1 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI2234

Product Information

Application	WB, FC
Primary Accession	Q16512
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	25GB1505
Calculated MW	103932
Gene Name	PKN1
Aliases	PKN1; Protein Kinase N1; PRK1; PAK1; PKN; Protein Kinase C-Like 1; PRKCL1; DBK; Serine/Threonine-Protein Kinase N1; Protease-Activated Kinase 1; Protein Kinase C-Like PKN; Protein Kinase PKN-Alpha; EC 2.7.11.13; MGC46204; PAK-1; Protein Kinase C-Related Kinase 1; Serine/Threonine Protein Kinase N; Protein-Kinase C-Related Kinase 1; Serine-Threonine Protein Kinase N; Serine-Threonine Kinase N; PKN-ALPHA; EC 2.7.11
Immunogen	A synthesized peptide derived from human PKN1

Additional Information

Gene ID	5585
Other Names	Serine/threonine-protein kinase N1, 2.7.11.13, Protease-activated kinase 1, PAK-1, Protein kinase C-like 1, Protein kinase C-like PKN, Protein kinase PKN-alpha, Protein-kinase C-related kinase 1, Serine-threonine protein kinase N, PKN1, PAK1, PKN, PRK1, PRKCL1

Protein Information

Name	PKN1
Synonyms	PAK1, PKN, PRK1, PRKCL1
Function	PKC-related serine/threonine-protein kinase involved in various processes such as regulation of the intermediate filaments of the actin cytoskeleton, cell migration, tumor cell invasion and transcription regulation. Part of a signaling cascade that begins with the activation of the adrenergic receptor ADRA1B and leads to the activation of MAPK14. Regulates the cytoskeletal network by phosphorylating proteins such as VIM and neurofilament proteins NEFH, NEFL and NEFM, leading to inhibit their polymerization. Phosphorylates 'Ser-575', 'Ser-637' and 'Ser-669' of MAPT/Tau, lowering its ability to bind to microtubules, resulting in disruption of tubulin assembly. Acts as a key coactivator of androgen receptor (AR)-dependent transcription, by being recruited to AR target genes and specifically mediating phosphorylation of

'Thr-11' of histone H3 (H3T11ph), a specific tag for epigenetic transcriptional activation that promotes demethylation of histone H3 'Lys-9' (H3K9me) by KDM4C/JMJD2C. Phosphorylates HDAC5, HDAC7 and HDAC9, leading to impair their import in the nucleus. Phosphorylates 'Thr-38' of PPP1R14A, 'Ser-159', 'Ser-163' and 'Ser-170' of MARCKS, and GFAP. Able to phosphorylate RPS6 in vitro.

Cellular Location

Cytoplasm. Nucleus Endosome. Cell membrane
{ECO:0000250|UniProtKB:Q63433}; Peripheral membrane protein
{ECO:0000250|UniProtKB:Q63433}. Cleavage furrow. Midbody
Note=Associates with chromatin in a ligand-dependent manner Localization to endosomes is mediated via its interaction with RHOB Association to the cell membrane is dependent on Ser-377 phosphorylation. Accumulates during telophase at the cleavage furrow and finally concentrates around the midbody in cytokinesis {ECO:0000250|UniProtKB:Q63433, ECO:0000269|PubMed:17332740}

Tissue Location

Found ubiquitously. Expressed in heart, brain, placenta, lung, skeletal muscle, kidney and pancreas. Expressed in numerous tumor cell lines, especially in breast tumor cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.