

KD-Validated Anti-Cullin 5 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

Catalog # AGI2249

Product Information

Application	WB
Primary Accession	Q93034
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Names	25GB4280
Calculated MW	90955
Gene Name	CUL5
Aliases	CUL5; Cullin 5; VACM-1; Vasopressin-Activated Calcium-Mobilizing Receptor 1; Cullin-5; CUL-5; VACM1; Cullin-5 (Vasopressin-Activated Calcium-Mobilizing Receptor-1); Vasopressin-Activated Calcium-Mobilizing Receptor-1
Immunogen	Recombinant protein of human Cullin 5

Additional Information

Gene ID	8065
Other Names	Cullin-5, CUL-5, Vasopressin-activated calcium-mobilizing receptor 1, VACM-1, CUL5 {ECO:0000303 PubMed:10230407, ECO:0000312 HGNC:HGNC:2556}

Protein Information

Name	CUL5 {ECO:0000303 PubMed:10230407, ECO:0000312 HGNC:HGNC:2556}
Function	Core component of multiple cullin-5-RING E3 ubiquitin-protein ligase complexes (ECS complexes, also named CRL5 complexes), which mediate the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:11384984, PubMed:15601820, PubMed:21199876, PubMed:21980433, PubMed:23897481, PubMed:25505247, PubMed:27910872, PubMed:32200094, PubMed:33268465, PubMed:35512830, PubMed:38418882, PubMed:40963025). Acts a scaffold protein that contributes to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme (PubMed:11384984, PubMed:15601820, PubMed:33268465). The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable SOCS box-containing substrate recognition component (PubMed:11384984, PubMed:15601820, PubMed:33268465). Acts as a key regulator of neuron positioning during cortex development: component of various SOCS-containing ECS complexes, such as the ECS(SOCS7) complex, that regulate reelin signaling by mediating ubiquitination and degradation of DAB1 (By similarity). ECS(SOCS1) seems to direct ubiquitination of JAK2 (PubMed:11384984). The ECS(SOCS2) complex mediates the ubiquitination and subsequent proteasomal degradation of

phosphorylated EPOR and GHR (PubMed:[21980433](#), PubMed:[25505247](#)). The ECS(PSB3) complex catalyzes ubiquitination of nuclear CGAS (PubMed:[38418882](#)). ECS(KLHDC1) complex is part of the DescEND (destruction via C-end degrons) pathway and mediates ubiquitination and degradation of truncated SELENOS selenoprotein produced by failed UGA/Sec decoding, which ends with a glycine (PubMed:[32200094](#)). The ECS(ASB9) complex mediates ubiquitination and degradation of CKB (PubMed:[33268465](#)). As part of some ECS complex, promotes 'Lys-11'-linked ubiquitination and degradation of BTRC (PubMed:[27910872](#)). As part of a multisubunit ECS complex, polyubiquitinates monoubiquitinated POLR2A (PubMed:[19920177](#)). As part of the ECS(RAB40C) complex, mediates ANKRD28 ubiquitination and degradation, thereby inhibiting protein phosphatase 6 (PP6) complex activity and focal adhesion assembly during cell migration (PubMed:[35512830](#)). As part of the ECS(RAB40A) complex, mediates RHOU 'Lys-48'-linked ubiquitination and degradation, thus inhibiting focal adhesion disassembly during cell migration (PubMed:[26598620](#)). As part of the ECS(RAB40B) complex, mediates LIMA1/EPLIN and RAP2 ubiquitination, thereby regulating actin cytoskeleton dynamics and stress fiber formation during cell migration (PubMed:[33999101](#), PubMed:[35293963](#)). The ECS(ASB7) complex acts a negative regulator of H3K9me3 histone mark by mediating ubiquitination and degradation of SUV39H1 (PubMed:[40440427](#)). As part of the ECS(LRRC58) complex, mediates CDO1 ubiquitination and degradation, regulating cysteine catabolism (PubMed:[40963025](#)). May form a cell surface vasopressin receptor (PubMed:[9037604](#)).

Cellular Location

Nucleus. Note=Localizes to sites of DNA damage in a UBAP2 and UBAP2L-dependent manner.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.